

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
17 February 2005 (17.02.2005)

PCT

(10) International Publication Number
WO 2005/014841 A1

(51) International Patent Classification⁷: **C12P 13/08**,
C12R 1/19 // (C12P 13/08, C12R 1:19)

(21) International Application Number:
PCT/EP2004/008389

(22) International Filing Date: 27 July 2004 (27.07.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
103 35 254.6 1 August 2003 (01.08.2003) DE
60/491,982 4 August 2003 (04.08.2003) US
10 2004 029 340.6 17 June 2004 (17.06.2004) DE

(71) Applicant (for all designated States except US): **DE-GUSSA AG** [DE/DE]; Bennigsenplatz 1, 40474 Düsseldorf (DE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **KRUSE, Daniela** [DE/DE]; Hermannstrasse 1, 33602 Bielefeld (DE). **HERMANN, Thomas** [DE/DE]; Zirkonstrasse 8, 33739 Bielefeld (DE). **RIEPING, Mechthild** [DE/DE]; Mönkebergstrasse 1, 33619 Bielefeld (DE). **THIERBACH, Georg** [DE/DE]; Gunststrasse 21, 33613 Bielefeld (DE).

(74) Common Representative: **DEGUSSA AG**; Intellectual Property Management, Patente und Marken, Standort Hanau, Postfach 13 45, 63403 Hanau (DE).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declaration under Rule 4.17:


— of inventorship (Rule 4.17(iv)) for US only

Published:

— with international search report
— before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: PROCESS FOR THE PREPARATION OF L-THREONINE



(57) Abstract: The invention relates to an improved, continuous cell-recycle process for the fermentative preparation of L-threonine using bacteria of the Enterobacteriaceae family which produce L-threonine.

WO 2005/014841 A1

Process for the Preparation of L-Threonine

Field of the Invention

The invention relates to an improved process for the fermentative preparation of L-threonine using bacteria of the Enterobacteriaceae family.

Background of the Invention

L-Threonine is used in animal nutrition, in human medicine and in the pharmaceuticals industry.

It is known that L-threonine can be prepared by fermentation of strains of the Enterobacteriaceae family, in particular *Escherichia coli*. Because of the great importance of this amino acid, work is constantly being undertaken to improve the preparation processes. Improvements to the process can relate to fermentation measures, such as e.g. stirring and supply of oxygen, or the composition of the nutrient media, such as e.g. the sugar concentration during the fermentation, or the working up to the product form, by e.g. ion exchange chromatography, or the intrinsic output properties, i.e. those of genetic origin, of the bacterium itself.

It is known that threonine can be prepared by fermentation of bacteria of the Enterobacteriaceae family, in particular *Escherichia coli*, in the batch process or fed batch process. In the batch process, all the nutrients are initially introduced directly at the start of the fermentation. In the fed batch process an additional nutrient medium is fed to the culture. This feed can start directly at the start of culturing or after a certain culturing time has elapsed, for example when a component introduced with the first nutrient medium initially introduced has been consumed. At the end of the fermentation, the complete contents of the fermenter are harvested and the threonine contained in the fermentation

broth is isolated and purified or otherwise processed. This process is described, for example, in the patent specifications US 5,538,873, EP-B-0593792 and WO 01/4525 and by Okamoto et al. (Bioscience, Biotechnology, and
5 Biochemistry 61 (11), 1877 - 1882, 1997).

Another process for the preparation of threonine using bacteria of the Enterobacteriaceae family, in particular Escherichia coli, is described in the patent specification US 6,562,601. It comprises initially culturing the
10 bacterium by the fed batch process, threonine becoming concentrated in the fermentation broth. At a desired point in time, a portion, i.e. 10 to 99%, of the fermentation broth contained in the fermenter is harvested. The remaining portion of the fermentation broth remains in the
15 fermenter. The fermentation broth remaining in the fermenting tank is topped up with nutrient medium and a further fermentation is carried out by the fed batch process. The cycle described is optionally carried out several times.

20 In another type of fermentation, continuous fermentation or chemostat culturing, a nutrient medium is fed to the culture continuously, while culture broth is removed continuously or semi-continuously, so that the volume of the culture broth in the fermenter remains approximately
25 constant. In theory, a continuous fermentation can be operated without limitation. Prolonging the fermentation has a significantly positive effect on the overall productivity of a fermenter (average amount of product produced per hour), since the influence of the time between
30 two fermentations on the overall productivity of a fermenter is reduced.

Industrial utilization of continuous processes for the preparation of amino acids is described in only isolated cases. The main reason for this lies in the degeneration of
35 the culture often observed. Amino acids are often produced

with genetically optimized microorganisms which have been selected on the basis of high substrate/product yields. In contrast to their precursors, they often have poorer growth properties here. During a long-lasting culturing, a
5 reversion to variants which produce less may occur. This leads to a loss in the productivity of the culture.

US 6,025,169 and FR 2669935 disclose that amino acids can be prepared by aerobic fed batch/continuous or continuous culturing with cell recycling of microorganisms.

10 The Abstract of JP62-289192 describes the improvement of a continuous process for the fermentative preparation of amino acids with *Corynebacterium glutamicum*. Improvements were achieved by varying the carbon concentration in the nutrient solution fed in, the stirrer speed, the redox
15 potential, the oxygen concentration and the intensity of the aeration.

JP62-289192 relates to a continuous process for the fermentative preparation of amino acids with bacteria which produce amino acids, in which the process can be improved
20 if the concentration of the source of carbon in the nutrient solution fed in continuously is above 10% and the redox potential in the culture is more than -200 mV. It is disclosed that the use of several fermenters can improve the utilization of the source of carbon if relatively large
25 amounts of the starting substances remain in the culture broth removed.

In JP60180598 (abstract) L-threonine is produced with *Brevibacterium lactofermentum* in batch fermentations. At the end of the fermentation the cells are filtered off and
30 employed in a new fermentation with fresh medium.

EP-A-139592 describes a fed batch process for the fermentative preparation of amino acids. In this, culture broth is removed from the reactor and the cells are

separated off by filtration, under growth limitation, and recycled into the process.

US 5,260,216 describes a continuous, fermentative process for the preparation of amino acid using bacteria with cell recycling. For separation of the cells, culture broth is removed from the reactor continuously. Before the bacteria cells are separated off and recycled into the process, air bubbles are separated off from the culture broth in order to increase the flow resistance of the liquids in the pipelines.

Kiyoshi Toda, 2003, Theoretical and Methodological Studies of Continuous Microbial Bioreactors, J. Gen. Appl. Microbiol., 49, 219-233, describes a continuous tryptophan process with cell recycling. Compared with a batch culture, it was possible to increase the cell concentration by a factor of 10 and the rate of production of tryptophan by a factor of 0.57 with this process.

Object of the Invention

It was the object of this invention to provide new measures for improved fermentative preparation of L-threonine.

Summary of the Invention

The invention provides a fermentation process, which comprises a procedure in which

- a) a bacterium of the Enterobacteriaceae family which produces L-threonine is inoculated and cultured in at least a first nutrient medium,
- b) at least a further nutrient medium or several further nutrient media is/are then fed continuously to the culture in one (F1) or several feed streams (F1+), the further nutrient medium or the further nutrient media comprising at least one source of

carbon, at least one source of nitrogen and at least one source of phosphorus, under conditions which allow the formation of L-threonine, and at the same time culture broth is removed from the culture according to the total of the removal stream F2 plus that of the cell-depleted removal stream F3 or according to the total of several removal streams (F2+) plus that of the cell-depleted removal stream F3, which substantially corresponds/correspond to the feed stream F1 or the total of the feed streams F1+, wherein

c) the cells are completely or partly separated off from the removal stream or streams F4 with a return flow ratio, R, of greater than 0 and less than or equal to 1 ($0 < R \leq 1$) by appropriate processes and are recycled into the culturing step b).

Description of the Invention

In this process, as shown in figure 1: F1 = a nutrient medium feed stream according to the invention; F1+ = several nutrient media feed streams according to the invention; F2 = a removal stream according to the invention; F2+ = several removal streams according to the invention; F3 = cell-depleted removal stream according to the invention (= F4 - F5); F4 = one or more removal streams for separating off cells; and F5 = the partly recycled removal stream F4 (= F4 - F3).

A "cell-depleted removal stream" is understood as meaning a removal stream from which at least some of the cells (= biomass) of the bacteria which secrete L-threonine have been removed.

According to the invention, the plant output of a fermenter which produces L-threonine can be increased by culturing by the batch process or fed batch process in the first step a)

described above, at least one additional nutrient medium being employed if the fed batch process is used. In the step b) described, at least one further nutrient medium or several further nutrient media are fed continuously to the culture in one or several feed streams and at the same time culture broth is removed from the culture with at least one or several removal streams, which substantially corresponds/correspond to the feed stream or the total of the feed streams.

- 10 During step b), in a step c) the cells or biomass contained in the removal stream or streams are completely or partly separated off from the removal stream or streams F4 by suitable processes with a return flow ratio, R , of greater than 0 and less than or equal to 1 ($0 < R \leq 1$) by
- 15 appropriate processes and are recycled into the culturing step b).

The term plant output is understood as meaning that in a plant, such as e.g. a fermenter, the weight or amount of a product, e.g. L-threonine, is prepared with a certain yield and with a certain rate or productivity or space/time yield. These parameters largely determine the costs or the profitability of a process.

The return flow ratio, R , is defined as the ratio of the cell-depleted removal stream according to the invention or the flow of the biomass-depleted permeate or sediment overflow (F3) to the total feed streams of the fermentation (F1 or F1+). At $R = 1$ the cells are completely retained and growth tends towards zero, while $R = 0$ represents normal chemostatic operation (Horst Chmiel, Bioprozesstechnik: Einführung in die Bioverfahrenstechnik [Bioprocess Technology: Introduction to Bioprocess Technology], Gustav Fischer Verlag, Stuttgart 1991).

According to the invention, the return flow ratio should be in the range of $0 < R \leq 1$, preferably between $0.1 < R \leq 1$,

$0.2 < R \leq 1$, $0.3 < R \leq 1$, $0.4 < R \leq 1$, $0.5 < R \leq 1$, $0.6 < R \leq 1$, $0.7 < R \leq 1$, $0.8 < R \leq 1$, $0.9 < R \leq 1$.

By the process according to the invention, the biomass concentrations are increased compared with a free culture system. That is to say, if the same microorganism is employed, according to the invention the biomass concentrations are increased by at least 5%, preferably at least 10%, at least 15%, at least 20%, at least 25%, at least 30%, at least 50%, at least 80%, at least 100%, at least 150%, or even by at least 200% or more. "Free culture system" means a process without cell retention, that is to say with a free overflow. The biomass concentration in a free culture system typically remains less than 50 g/l, less than 40 g/l, less than 30 g/l, or even less than 20 g/l. These different biomass concentrations in a free culture system depend on the microorganisms employed and the processes employed.

Centrifuges, separators and settlers (after flocculation) as well as cross-stream, micro- and ultra-filters are suitable for separating off the cells (Biotechnologie [Biotechnology], by H. Weide, J. Páca and W. A. Knorre, Gustav Fischer Verlag, Jena, 1991). The separating off of cells can take place here in one step or by several separation steps, which, for example, can be arranged sequentially. The biomass which has been separated off is then recycled into the fermenter.

A culture broth is understood as meaning the suspension of a microorganism formed by culturing a microorganism - in the case of the present invention a bacterium which produces L-threonine - in a nutrient medium using a fermenter or culture vessel.

During the culturing step a), the bacterium is inoculated in at least a first nutrient medium and cultured by the batch process or fed batch process. If the fed batch

process is used, an additional nutrient medium is fed in after more than 0 to not more than 10 hours, preferably after 1 to 10 hours, preferentially after 2 to 10 hours and particularly preferably after 3 to 7 hours.

- 5 The first nutrient medium comprises as the source of carbon one or more of the compounds chosen from the group consisting of sucrose, molasses from sugar beet or cane sugar, fructose, glucose, starch hydrolysate, lactose, galactose, maltose, xylose, cellulose hydrolysate,
10 arabinose, acetic acid, ethanol and methanol, in concentrations of 1 to 100 g/kg or 1 to 50 g/kg, preferably 10 to 45 g/kg, particularly preferably 20 to 40 g/kg. Starch hydrolysate is understood according to the invention as the hydrolysis product of starch from maize, cereals,
15 potatoes or tapioca.

- Organic nitrogen-containing compounds, such as peptones, yeast extract, meat extract, malt extract, corn steep liquor, soya bean flour and urea, or inorganic compounds, such as ammonia, ammonium sulfate, ammonium chloride,
20 ammonium phosphate, ammonium carbonate and ammonium nitrate, potassium nitrate and potassium sodium nitrate, can be used as the source of nitrogen in the first nutrient medium. The sources of nitrogen can be used individually or as a mixture in concentrations of 1 to 40 g/kg, preferably
25 10 to 30 g/kg, particularly preferably 10 to 25 g/kg, very particularly preferably 1 to 30 g/kg or 1 to 25 g/kg.

- Phosphoric acid, alkali metal or alkaline earth metal salts of phosphoric acid, in particular potassium dihydrogen phosphate or dipotassium hydrogen phosphate or the
30 corresponding sodium-containing salts, polymers of phosphoric acid or the hexaphosphoric acid ester of inositol, also called phytic acid, can be used as the source of phosphorus in the first nutrient medium in concentrations of 0.1 to 5 g/kg, preferably 0.3 to 3 g/kg,
35 particularly preferably 0.5 to 1.5 g/kg. The culture medium

must furthermore comprise salts of metals, such as e.g. magnesium sulfate or iron sulfate, which are necessary for growth. These substances are present in concentrations of 0.003 to 3 g/kg. Finally, essential growth substances, such as amino acids (e.g. homoserine) and vitamins (e.g. thiamine), are employed in addition to the above-mentioned substances. Antifoams, such as e.g. fatty acid polyglycol esters, can be employed to control the development of foam.

The additional nutrient medium which is used in a fed batch process in general comprises merely as the source of carbon one or more of the compounds chosen from the group consisting of sucrose, molasses from sugar beet or cane sugar, fructose, glucose, starch hydrolysate, lactose, galactose, maltose, xylose, cellulose hydrolysate, arabinose, acetic acid, ethanol and methanol, in concentrations of 300 to 700 g/kg, preferably 400 to 650 g/kg, and optionally an inorganic source of nitrogen, such as e.g. ammonia, ammonium sulfate, ammonium chloride, ammonium phosphate, ammonium carbonate, ammonium nitrate, potassium nitrate or potassium sodium nitrate. Alternatively, these and other components can also be fed in separately.

It has been found that in the continuous culturing according to step b), the constituents of the further nutrient medium can be fed to the culture in the form of a single further nutrient medium and in a plurality of further nutrient media. According to the invention, the further nutrient medium or the further nutrient media are fed to the culture in at least one (1) feed stream or in a plurality of feed streams of at least 2 to 10, preferably 2 to 7 or 2 to 5 feed streams.

The term "continuous" means that the feed stream or the feed streams are substantially uninterrupted. That is to say, nutrient medium or nutrient media is/are added to the culture with at most short, individual pauses. The

individual interruptions or pauses are up to a maximum of 0.5, a maximum of 1, a maximum of 2 or a maximum of 3 hours. The sum of the individual interruptions or pauses in the culturing according to step b) is a maximum of 10%,
5 a maximum of 8%, a maximum of 6%, a maximum of 4%, a maximum of 2% or a maximum of 1% of the total time of the culturing according to step b).

The further nutrient medium or the further nutrient media comprises/comprise as the source of carbon one or more of
10 the compounds chosen from the group consisting of sucrose, molasses from sugar beet or cane sugar, fructose, glucose, starch hydrolysate, maltose, xylose, cellulose hydrolysate, arabinose, acetic acid, ethanol and methanol, in concentrations of 20 to 700 g/kg, preferably 50 to
15 650 g/kg.

The further nutrient medium or the further nutrient media furthermore comprises or comprise a source of nitrogen consisting of organic nitrogen-containing compounds, such as peptones, yeast extract, meat extract, malt extract,
20 corn steep liquor, soya bean flour and urea, or inorganic compounds, such as ammonia, ammonium sulfate, ammonium chloride, ammonium phosphate, ammonium carbonate, ammonium nitrate and/or potassium nitrate or potassium sodium nitrate. The sources of nitrogen can be used individually
25 or as a mixture in concentrations of 5 to 50 g/kg, preferably 10 to 40 g/kg.

The further nutrient medium or the further nutrient media furthermore comprises or comprise a source of phosphorus consisting of phosphoric acid or the alkali metal or
30 alkaline earth metal salts of phosphoric acid, in particular potassium dihydrogen phosphate or dipotassium hydrogen phosphate or the corresponding sodium-containing salts, polymers of phosphoric acid or the hexaphosphoric acid ester of inositol, also called phytic acid, or the
35 corresponding alkali metal or alkaline earth metal salts.

The sources of phosphorus can be used individually or as a mixture in concentrations of 0.3 to 3 g/kg, preferably 0.5 to 2 g/kg. The further nutrient medium or the further nutrient media must furthermore comprise salts of metals, 5 such as e.g. magnesium sulfate or iron sulfate, which are necessary for growth, in concentrations of 0.003 to 3 g/kg, preferably in concentrations of 0.008 to 2 g/kg. Finally, essential growth substances, such as amino acids (e.g. homoserine) and vitamins (e.g. thiamine), are employed in 10 addition to the above-mentioned substances. Antifoams, such as e.g. fatty acid polyglycol esters, can be employed to control the development of foam.

If a single further nutrient medium is used, this is typically fed to the culture in one feed stream. If a 15 plurality of further nutrient media are used, these are fed in a corresponding plurality of feed streams. If a plurality of further nutrient media are used, it should be noted that these in each case can comprise only one of the sources of carbon, nitrogen or phosphorus described, or 20 also a mixture of the sources of carbon, nitrogen or phosphorus described.

According to the invention, the nutrient medium fed in or the nutrient media fed in is/are adjusted such that there is a phosphorus to carbon ratio (P/C ratio) of not more 25 than 4; of not more than 3; of not more than 2; of not more than 1.5; of not more than 1; of not more than 0.7; of not more than 0.5; of not more than 0.48; of not more than 0.46; of not more than 0.44; of not more than 0.42; of not more than 0.40; of not more than 0.38; of not more than 30 0.36; of not more than 0.34; of not more than 0.32; or of not more than 0.30 mmol of phosphorus / mol of carbon.

The feed stream or the total of the feed streams in the process according to the invention are fed in at a rate corresponding to an average residence time of less than 30 35 hours, preferably less than 25, very particularly

preferably less than 20 hours. The average residence time here is the theoretical time the particles remain in a continuously operated culture. The average residence time is described by the ratio of the volume of liquid in the reactor and the amount flowing through (Biotechnologie [Biotechnology]; H. Weide, J. Páca and W. A. Knorre; Gustav Fischer Verlag Jena; 1991).

Intensive growth at the start of culturing according to step (a) is usually a logarithmic growth phase. The logarithmic growth phase is in general followed by a phase of less intensive cell growth than in the logarithmic phase.

After 10 to 72 hours, preferably 15 to 48 hours, or during or after the logarithmic growth phase, as described above at least one further nutrient medium or several further nutrient media are fed continuously to the culture in one or several feed streams and at the same time culture broth is removed from the culture with one or several removal streams, which substantially corresponds/respond to the feed stream or the total of the feed streams, the cells being completely or partly separated off from the removal stream or streams F4 and recycled into the culturing step b) with a return flow ratio, R, of greater than 0 and less than or equal to 1 ($0 < R \leq 1$).

Substantially here means that the total of the removal stream F2 plus F3 or the total of the removal streams F2+ plus F3 corresponds to 80% - 120%, 90% - 110% or 95% - 105% of the feed stream F1 or of the total of the feed streams F1+. The removal can be realized industrially by pumping off and/or by draining off the culture broth.

According to the invention, the concentration of the source of carbon during the culturing according to step b) and/or c) is in general adjusted to not more than 30 g/l, to not more than 20 g/l, to not more than 10 g/l, preferably to

not more than 5 g/l, particularly preferably to not more than 2 g/l. This concentration is maintained for at least 75%, preferably for at least 85%, particularly preferably for at least 95% of the time of the culturing according to
5 step b) and/or c). The concentration of the source of carbon is determined here with the aid of methods which are prior art. β -D-Glucose is determined e.g. in a YSI 02700 Select glucose analyzer from Yellow Springs Instruments (Yellow Springs, Ohio, USA).

10 Brief Description of the Figure

Figure 1 is a diagram which shows the feed and the removal and return flow streams of a process according to the invention. In this figure 1 the symbols denote the following:

15 F1 = a nutrient medium feed stream according to the invention; F1+ = several nutrient media feed streams according to the invention; F2 = a removal stream according to the invention; F2+ = several removal streams according to the invention; F3 = cell-depleted removal stream
20 according to the invention ($= F4 - F5$); F4 = one or several removal streams for separating off cells; and F5 = the partly recycled removal stream F4 ($= F4 - F3$).

As figure 1 shows, according to the invention cells are completely or partly separated off from the removal stream
25 (F4) with a return flow ratio, R, of $0 < R \leq 1$ by appropriate processes and are recycled as F5 into the culturing step (b). This return flow ratio is equivalent to a recycling of greater than 0% up to and including 100% of the removed cells or biomass, preferably of 10% up to and
30 including 100%, of 20% up to and including 100%, of 30% up to and including 100%, of 40% up to and including 100%, of 50% up to and including 100%, of 60% up to and including 100%, of 70% up to and including 100%, of 80% up to and including 100%, or of 90% up to and including 100%.

- The separating off of cells can take place here e.g. by micro- or ultrafiltration in one step, or sequentially by various centrifugation or separation steps. If the separating off of the biomass takes place in one step to
5 100%, some of the biomass retained can be recycled back into the culture vessel/the fermenter, and in particular corresponding to a return flow ratio R of $0 < R \leq 1$. In the case of sequential separating off of the biomass, for example by sequential centrifugation steps or sequential
10 sedimentation steps, the return flow ratio R can be determined by a particular fraction of the centrifugate being recycled. Particular fractions can furthermore be mixed with one another in order to establish a desired return flow ratio.
- 15 The return flow ratio is adjusted by determining the biomass concentration in the culture vessel or fermenter and/or determining the biomass concentration in the return flow. The biomass concentration is determined by the techniques conventionally used, such as e.g. counting the
20 cell titer in a calibrated counting chamber, cytometrically with or without staining of the cells, determination of the optical density of the culture or by determination of the dry biomass. However, counting of the cell titer can also be automated, for example by determining the biomass
25 concentration in a filtrate or centrifugate stage by optical methods, such as photometry or cytometry, or by using physical methods, such as, for example, determination of the conductivity or the absorption of radiation of the near infra-red or middle infra-red range.
- 30 If appropriate, the culture broth removed can be provided with oxygen or an oxygen-containing gas until the concentration of the source of carbon falls below 2 g/l; below 1 g/l; or below 0.5 g/l.

In a process according to the invention, the yield is at
35 least 31%, at least 33%, at least 35%, at least 37%, at

least 38%, at least 40%, at least 42%, at least 44%, at least 46% or at least 48%. The yield is defined here as the ratio of the total amount of L-threonine formed in a culturing to the total amount of the source of carbon employed or consumed.

In a process according to the invention, L-threonine is formed with a space/time yield of at least 1.5 to 2.5 g/l per h, of at least 2.5 to 3.5 g/l per h, of at least 2.5 to more than 3.5 g/l per h, of at least 3.5 to 5.0 g/l per h, of at least 3.5 to more than 5.0 g/l per h, or of at least 5.0 to 8.0 g/l per h or more, such as, for example, at least 9 g/l per h. The space/time yield is defined here as the ratio of the total amount of threonine formed in a culturing to the volume of the culture over the total period of time of culturing. The space/time yield is also called the volumetric productivity.

In a fermentation process like that according to the invention, the product is of course prepared in a certain yield and in a certain space/time yield (volumetric productivity). In a process according to the invention, L-threonine can be prepared in a yield of at least 31% and a space/time yield of at least 1.5 to 2.5 g/l per h. Further couplings of yield with space/time yield, such as, for example, a yield of at least 37% and a space/time yield of at least 2.5 g/l per h, automatically result from the above statements.

The culturing in steps a) and b) is carried out under conditions which allow the formation of L-threonine: During the culturing the temperature is adjusted in a range from 29 to 42°C, preferably 33 to 40°C. The culturing can be carried out under normal pressure or optionally under increased pressure, preferably under an increased pressure of 0 to 1.5 bar. The oxygen partial pressure is regulated at 5 to 50%, preferably approx. 20% atmospheric saturation. During this procedure the culture is stirred and supplied

with oxygen. Regulation of the pH to a pH of approx. 6 to 8, preferably 6.5 to 7.5, can be effected with 25% aqueous ammonia.

The process according to the invention is operated for at least approx. 72 hours, preferably 100 to ≥ 300 hours, particularly preferably 200 to ≥ 300 hours. In the process according to the invention, the volume of the culture is exchanged at least by half, at least once, at least 2 times, at least 3 times, at least 4 times, at least 6 times, at least 8 times, at least 10 times, at least 12 times.

From the culture broth removed, the L-threonine can be isolated, collected or concentrated and optionally purified.

It is also possible to prepare a product from the culture broth removed (= fermentation broth) by removing the bacterium biomass contained in the culture broth completely (100%) or almost completely, i.e. more than or greater than (>) 90%, >95%, >97%, >99% and leaving the other constituents of the fermentation broth largely, i.e. to the extent of 30% - 100%, 40% - 100%, 50% - 100%, 60% - 100%, 70% - 100%, 80% - 100%, or 90% - 100%, preferably greater than or equal to (\geq) 50%, $\geq 60\%$, $\geq 70\%$, $\geq 80\%$, $\geq 90\%$ or $\geq 95\%$ or also completely (100%) in the product.

Separation methods such as, for example, centrifugation, filtration, decanting, flocculation or a combination thereof are employed for the removal or separating off of the biomass.

The broth obtained is then thickened or concentrated by known methods, such as, for example, with the aid of a rotary evaporator, thin film evaporator, falling film evaporator, by reverse osmosis, by nanofiltration or a combination thereof.

This concentrated broth is then be worked up by methods of freeze drying, spray drying, spray granulation or by other processes to give a preferably free-flowing, finely divided powder. This free-flowing, finely divided powder can then
5 in turn by converted by suitable compacting or granulating processes into a coarse-grained, readily free-flowing, storable and largely dust-free product. The water is removed in total to the extent of more than 90% by this means, so that the water content in the product is less
10 than 10%, less than 5%.

The process steps mentioned do not necessarily have to be carried out in the sequence stated here, but can optionally be combined in an industrially appropriate manner.

The analysis of L-threonine and other amino acids can be
15 carried out by anion exchange chromatography with subsequent ninhydrin derivation, as described by Spackman et al. (Analytical Chemistry, 30: 1190-1206 (1958)) or it can be carried out by reversed phase HPLC, as described by Lindroth et al. (Analytical Chemistry 51: 1167-1174
20 (1979)).

Bacteria of the Enterobacteriaceae family which produce L-threonine chosen from the genera Escherichia, Erwinia, Providencia and Serratia are suitable for carrying out the process according to the invention. The genera Escherichia
25 and Serratia are preferred. Of the genus Escherichia in particular the species Escherichia coli and of the genus Serratia in particular the species Serratia marcescens are to be mentioned.

The bacteria contain at least one copy of a thrA gene or
30 allele which codes for a threonine-insensitive aspartate kinase I - homoserine dehydrogenase I. In this connection, "feed back" resistant or also desensitized variants are referred to in the literature. Such bacteria are typically resistant to the threonine analogue α -amino- β -

hydroxyvaleric acid (AHV) (Shiio and Nakamori, Agricultural and Biological Chemistry 33 (8), 1152-1160 (1969)).

Biochemical studies on "feed back" resistant aspartate kinase I - homoserine dehydrogenase I variants are

5 described, for example, by Cohen et al. (Biochemical and Biophysical Research Communications 19(4), 546-550 (1965)) and by Omori et al. (Journal of Bacteriology 175(3), 785-794 (1993)). If appropriate, the threonine-insensitive aspartate kinase I - homoserine dehydrogenase I is
10 overexpressed.

Methods of overexpression are adequately described in the prior art - for example by Makrides et al. (Microbiological Reviews 60 (3), 512-538 (1996)). By using vectors, the number of copies is increased by at least one (1) copy.

15 Vectors which can be used are plasmids such as are described, for example, in US 5,538,873. Vectors which can also be used are phages, for example the phage Mu, as described in EP 0 332 448, or the phage lambda (λ). An increase in the number of copies can also be achieved by
20 incorporating a further copy into a further site of the chromosome - for example into the att site of the phage λ (Yu and Court, Gene 223, 77-81 (1998)). US 5,939,307 discloses that by incorporation of expression cassettes or promoters, such as, for example, the tac promoter, trp
25 promoter, lpp promoter or P_L promoter and P_R -promoter of the phage λ , upstream of the chromosomal threonine operon it was possible to achieve an increase in the expression. The promoters of the phage T7, the gear-box promoters or the nar promoter can be used in the same way. Such expression
30 cassettes or promoters can also be used, as described in EP 0 593 792, to overexpress plasmid-bound genes. By using the $lacI^Q$ allele, the expression of plasmid-bound genes can in turn be controlled (Glascock and Weickert, Gene 223, 221-231 (1998)). By removal of the attenuator of the
35 threonine operon (Park et al., Biotechnology Letters 24, 1815-1819 (2002)) or by using the thr79-20 mutation

(Gardner, Proceedings of the National Academy of Sciences, USA 76(4), 1706-1710 (1979)) or by mutation of the thrS gene, which codes for threonyl-t-RNA synthetase, as described by Johnson et al. (Journal of Bacteriology 5 129(1), 66-70 (1977) an overexpression can likewise be achieved. By the measures described, the intracellular concentration of the particular aspartate kinase I - homoserine dehydrogenase I protein variant is increased by at least 10% compared with the starting strain.

10 A suitable thrA-allele is described in US 4,278,765 and is obtainable in the form of the strain MG442 from the Russische Nationalsammlung für industrielle Mikroorganismen [Russian National Collection of Industrial Microorganisms] (VKPM, Moscow, Russia) under the Accession Number CMIM 15 B-1628. Other suitable thrA alleles are described in WO 00/09660 and WO 00/09661 and are obtainable from the Korean Culture Center of Microorganisms (KCCM, Seoul, Korea) under the Accession Numbers KCCM 10132 and KCCM 10133. A further suitable thrA allele is present in the 20 strain H-4581, which is described in US 4,996,147 and is obtainable under the Accession Number Ferm BP-1411 from the National Institute of Advanced Industrial Science and Technology (1-1-1 Higashi, Tsukuba Ibaraki, Japan). Finally, further thrA alleles are described in US 3,580,810 25 and are obtainable in the form of the strains ATCC 21277 and ATCC 21278 deposited at the ATCC. A further allele is described in US 3,622,453 and is obtainable from the ATCC in the form of the strain KY8284 under the Accession Number ATCC 21272. Furthermore, WO 02/064808 describes a further 30 thrA allele which is deposited at the KCCM in the form of strain pGmTN-PPC12 under the Accession Number KCCM 10236.

If appropriate, thrA alleles which code for "feed back" resistant aspartate kinase I - homoserine dehydrogenase I variants can be isolated using the adequately known methods 35 of conventional mutagenesis of cells using mutagenic

substances, for example N-methyl-N'-nitro-N-nitrosoguanidine (MNNG) or ethyl methanesulfonate (EMS) or mutagenic rays, for example UV rays, and subsequent selection of threonine analogues (for example AHV-resistant variants). Such mutagenesis methods are described, for example, by Shiio and Nakamori (Agricultural and Biological Chemistry 33 (8), 1152-1160 (1969)) or by Saint-Girons and Margerita (Molecular and General Genetics 162, 101-107 (1978)) or in the known handbook of J. H. Miller (A Short Course In Bacterial Genetics. A Laboratory Manual and Handbook for Escherichia coli and Related Bacteria, Cold Spring Harbor Laboratory Press, New York, USA, 1992), in particular on pages 135 to 156. Shiio and Nakamori, for example, treat a cell suspension of Escherichia coli with 0.5 mg/ml MNNG in a 0.1 M sodium phosphate buffer of pH 7 at room temperature (i.e. in general approx. 16 to 26°C) for approx. 15 minutes to generate mutations. Miller recommends, for example, a treatment for 5 to 60 minutes with 30 µl EMS per 2 ml of cell suspension in 0.1 M TRIS buffer at pH 7.5 at a temperature of 37°C. These mutagenesis conditions can be modified in an obvious manner. Selection of AHV-resistant mutants takes place on minimal agar, which typically contains 2 to 10 mM AHV. The corresponding alleles can then be cloned and subjected to a sequence determination and the protein variants coded by these alleles can be subjected to a determination of the activity. If appropriate, the mutants produced can also be used directly. The word "directly" means that the mutants produced can be employed for the preparation of L-threonine in a process according to the invention or that further modifications can be carried out on these mutants to increase the output properties, such as, for example, attenuation of the threonine degradation or overexpression of the threonine operon.

Methods of in vitro mutagenesis such as are described, for example, in the known handbook by Sambrook et al.

(Molecular Cloning, A Laboratory Manual, 2nd ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York, USA, 1989) can also be used in the same way.

Corresponding methods are also available commercially in the form of so-called "kits", such as, for example, the "QuikChange Site-Directed Mutagenesis Kit" from Stratagene (La Jolla, USA) described by Papworth et al. (Strategies 9(3), 3-4 (1996)).

These mutagenesis methods can of course also be used on other genes, alleles or strain or objectives and tasks, such as, for example, the production and isolation of mutants which are resistant to L-threonine.

Those thrA alleles which code for aspartate kinase I - homoserine dehydrogenase I variants which, in the presence of 10 mM L-threonine, have at least 40%, at least 45%, at least 50%, at least 55% or at least 60% of the homoserine dehydrogenase activity and/or which, in the presence of 1 mM L-threonine, have at least 60%, at least 70%, at least 75% or at least 80% of the homoserine dehydrogenase activity, compared with the activity in the absence of L-threonine, are preferred. Where appropriate, the aspartate kinase activity of the aspartate kinase I - homoserine dehydrogenase I variants mentioned in the presence of 10 mM L-threonine is at least 60%, at least 65%, at least 70%, at least 75% or at least 80% of the activity in the absence of L-threonine.

Bacteria of the Enterobacteriaceae family which contain a stop codon chosen from the group consisting of opal, ochre and amber, preferably amber in the rpoS gene, and a t-RNA suppressor chosen from the group consisting of opal suppressor, ochre suppressor and amber suppressor, preferably amber suppressor, are moreover suitable. The amber mutation preferably lies at position 33 according to the amino acid sequence of the RpoS gene product. supE is preferably employed as the amber suppressor. These bacteria

are described in PCT/EP02/02055. A strain which contains the mutation described in the *rpoS* gene and the suppressor *supE* is obtainable under the Accession Number DSM 15189 from the Deutsche Sammlung für Mikroorganismen und
5 Zellkulturen [German Collection of Microorganisms and Cell Cultures] (Braunschweig, Germany).

The nucleotide sequence of the *rpoS* gene can be found in the prior art. The nucleotide sequence of the *rpoS* gene corresponding to Accession No. AE000358 is shown as SEQ ID
10 NO. 1. The amino acid sequence of the associated RpoS gene product or protein is shown in SEQ ID NO. 2. The nucleotide sequence of an *rpoS* allele which contains a stop codon of the amber type at the position of the nucleotide sequence corresponding to position 33 of the amino acid sequence of
15 the RpoS gene product or protein, corresponding to SEQ ID NO. 1 or SEQ ID NO. 2 respectively, is reproduced in SEQ ID NO. 3. The suppressor *supE* is described in the prior art and is shown as SEQ ID NO. 4.

Bacteria of the Enterobacteriaceae family which are not
20 capable of breaking down threonine or utilizing it as a source of nitrogen under aerobic culture conditions are moreover suitable. Aerobic culture conditions are understood as meaning those under which the oxygen partial pressure in the fermentation culture is greater than (>) 0%
25 during 90%, preferably 95%, very particularly preferably 99% of the duration of the fermentation. Such a strain is, for example, the strain KY10935 described by Okamoto (Bioscience, Biotechnology and Biochemistry 61(11), 1877-1882 (1997)). Strains which are not capable of breaking
30 down threonine, with splitting off of nitrogen, in general have an attenuated threonine dehydrogenase (EC 1.1.1.103), which is coded by the *tdh* gene. The enzyme has been described by Aronson et al. (The Journal of Biological Chemistry 264(9), 5226-5232 (1989)). Attenuated *tdh* genes
35 are described, for example, by Ravnkar and Somerville

(Journal of Bacteriology, 1986, 168(1), 434-436), in US 5,705,371, in WO 02/26993 and by Komatsubara (Bioprocess Technology 19, 467-484 (1994)).

- A suitable tdh allele is described in US 5,538,873 and is
5 obtainable in the form of the strain B-3996 under the
Accession Number 1876 from the Russische Nationalsammlung
für industrielle Mikroorganismen [Russian National
Collection of Industrial Microorganisms] (VKPM, Moscow,
Russia). A further tdh allele is described in US 5,939,307
10 and is obtainable in the form of the strain kat-13 under
the Accession Number NRRL B-21593 from the Agriculture
Research Service Patent Culture Collection (Peoria,
Illinois, USA). Finally, a tdh allele is described in WO
02/26993 and is deposited in the form of the strain TH21.97
15 under the Accession Number NRRL B-30318 at the NRRL. The
allele tdh-1::cat1212, which codes for a defective
threonine dehydrogenase, is obtainable from the E. coli
Genetic Stock Center (New Haven, Conn., USA) under the
Accession Number CGSC 6945.
- 20 Bacteria of the Enterobacteriaceae family which have an at
least partial need for isoleucine ("leaky phenotype") which
can be compensated by addition of L-isoleucine in a
concentration of at least 10, 20 or 50 mg/l or L-threonine
in a concentration of at least 50, 100 or 500 mg/l are
25 moreover suitable.

- Need or auxotrophy is in general understood as meaning the
fact that as a result of a mutation, a strain has
completely lost a wild-type function, for example an enzyme
activity, and requires the addition of a supplement, for
30 example an amino acid, for growth. A partial need or
partial auxotrophy is referred to if, as a result of a
mutation, a wild-type function, for example the activity of
an enzyme from the biosynthesis pathway of an amino acid,
is impaired or attenuated but not eliminated completely.
35 Strains with a partial need typically have, in the absence

of the supplement, a growth rate which is reduced, i.e. greater than (>) 0% and less than (<) 90%, 50%, 25% or 10%, compared with the wild-type. In the literature, this relationship is also called "leaky" phenotype or

- 5 "leakiness" (Griffiths et al.: An Introduction to Genetic Analysis. 6th edition, 1996, Freeman and Company, New York, USA).

A strain with such a partial need for isoleucine is described, for example, in WO 01/14525 and deposited in the
10 form of the strain DSM9906 under the Accession Number KCCM 10168 at the KCCM. Threonine-secreting or -producing strains with a need for isoleucine in general have an attenuated threonine deaminase (E.C. Number 4.3.1.19), which is coded by the *ilvA* gene. Threonine deaminase is
15 also known by the name threonine dehydratase. An attenuated *ilvA* gene which effects a partial isoleucine auxotrophy is described, for example, in US 4,278,765 and is obtainable in the form of the strain MG442, deposited under the Accession Number B-1682, at the VKPM.

- 20 A further attenuated *ilvA* gene is described, for example, in WO 00/09660 and deposited in the form of the strain DSM 9807 under the Accession Number KCCM -10132 at the KCCM. Further attenuated *ilvA* genes are described by Komatsubara (Bioprocess Technology 19, 467-484 (1994)).

- 25 The amino acid sequence of a suitable and new threonine deaminase comprises, for example, the sequence of SEQ ID NO. 6, which can contain any amino acid apart from glutamic acid at position 286. Exchange of glutamic acid for lysine is preferred (E286K).

- 30 The term "amino acid" means, in particular, the proteinogenic L-amino acids, including their salts, chosen from the group consisting of L-asparagine, L-threonine, L-serine, L-glutamate, L-glycine, L-alanine, L-cysteine, L-valine, L-methionine, L-isoleucine, L-leucine, L-tyrosine,

L-phenylalanine, L-histidine, L-lysine, L-tryptophan, L-proline and L-arginine.

SEQ ID NO. 8 shows the amino acid sequence of a threonine deaminase which contains the amino acid lysine at position 286; the associated nucleotide sequence is shown as SEQ ID NO. 7. This contains the nucleobase adenine at position 856.

Another suitable threonine deaminase is the variant described by Lee et al. (Journal of Bacteriology 185 (18), 5442-5451 (2003)), in which serine is exchanged for phenylalanine at position 97 (S97F). Further suitable threonine deaminases are the variants described by Fischer and Eisenstein (Journal of Bacteriology 175 (20), 6605-6613 (1993)), which have at least one of the amino acid exchanges chosen from the group consisting of: exchange of asparagine at position 46 for aspartic acid (N46D), exchange of alanine at position 66 for valine (A66V), exchange of proline at position 156 for serine (P156S), exchange of glycine at position 248 for cysteine (G248C) and exchange of aspartic acid at position 266 for tyrosine (D266Y).

By insertion or deletion mutagenesis of at least one base pair or nucleotide or by insertion or deletion of at least one codon in the coding region or by incorporation of a stop codon by transition or transversion mutagenesis into the coding region of the *ilvA* gene, alleles in which the expression of the *ilvA* gene is in general completely eliminated can be isolated. This method can also be applied to other genes, alleles or open reading frames, such as, for example, the *tdh* gene, which codes for threonine dehydrogenase.

Bacteria of the Enterobacteriaceae family which are resistant in their growth towards inhibition by L-threonine and/or L-homoserine are moreover suitable. Threonine-

resistant strains and the preparation thereof are described, for example, by Astaurova et al. (Prikladnaya Biokhimiya Microbiologiya (1985), 21(5), 485 as the English translation: Applied Biochemistry and Microbiology (1986), 21, 485-490)). The mutant described by Austaurova is resistant towards 40 mg/ml L-threonine. Furthermore, for example, the strain 472T23, which can grow in the presence of 5 mg/ml L-threonine and at the same time is resistant to L-homoserine, is described in US 5,175,107. The strain 472T232 is obtainable under the Accession Number BKIIM B-2307 from the VKPM and under the number ATCC 9801 from the ATCC. Furthermore, the strain DSM 9807, which can grow on a solid nutrient medium which comprises 7% L-threonine, is described in WO 00/09660. The strain DSM 9807 is obtainable under the Accession Number KCCM-10132 from the KCCM. Finally, the strain DSM 9906, which can grow in a medium which comprises 60% to 70% of an L-threonine fermentation mother liquid, is described in WO 01/14525. The strain DSM 9906 is obtainable under the Accession Number KCCM-10168 from the KCCM.

It is known (see EP 0994 190 A2 and Livshits et al. (Research in Microbiology 154, 123-135 (2003)), that resistance to L-threonine and L-homoserine is brought about by enhancement of the rhtA gene. The enhancement can be achieved by increasing the number of copies of the gene or by use of the rhtA23 mutation.

EP 0 994 190 A2 discloses that the enhancement of the rhtB gene effects resistance to L-homoserine and L-threonine, in particular to L-homoserine, and improves threonine production. By overexpression of the RhtB gene product in a strain called N99, it was possible to increase the minimum inhibitory concentration from 250 µg/ml to 30,000 µg/ml.

EP 1,013,765 A1 discloses that an enhancement of the rhtC gene causes resistance to L-threonine and improves threonine production. A strain which can grow on a minimal

agar in the presence of a concentration of at least 30 mg/ml L-threonine is called resistant to L-threonine. It is furthermore disclosed that an enhancement of the *rhtB* gene effects resistance to L-homoserine and improves

5 threonine production. A strain which can grow on a minimal agar in the presence of a concentration of at least 5 mg/ml L-homoserine is called resistant to L-homoserine. The patent application mentioned describes strains which are resistant to 10 mg/ml L-homoserine and resistant to 50

10 mg/ml L-threonine. US 4,996,147 describes the strain H-4581, which is resistant to 15 g/l homoserine. The strain H-4581 is obtainable under the Accession Number FERM BP-1411 from the National Institute of Advanced Industrial Science and Technology.

15 EP 1 016 710 A2 discloses that an enhancement of the open reading frame or gene *yfiK* or *yeaS* effects resistance to L-threonine and L-homoserine. By overexpression of the *YfiK* gene product in a strain called TG1, it was possible to increase the minimum inhibitory concentration in respect of

20 L-homoserine from 500 µg/ml to 1,000 µg/ml and in respect of L-threonine from 30,000 µg/ml to 40,000 µg/ml. By overexpression of the *YeaS* gene product, it was possible to increase the minimum inhibitory concentration in respect of L-homoserine from 500 µg/ml to 1,000 µg/ml and in respect

25 of L-threonine from 30,000 µg/ml to 50,000 µg/ml. It is furthermore demonstrated in the patent application mentioned that threonine production is improved by overexpression of the *YfiK* gene product.

Strains which can grow in the presence of \geq (at least)

30 5 g/l, ≥ 10 , ≥ 20 g/l, ≥ 30 g/l, ≥ 40 g/l, ≥ 50 g/l, ≥ 60 g/l and ≥ 70 g/l L-threonine, i.e. are resistant to L-threonine, and are suitable for the preparation of L-threonine in a process according to the invention are prepared according to these technical instructions.

Strains which have at least the following features are suitable in particular for the process according to the invention:

- 5 a) a threonine-insensitive aspartate kinase I - homoserine dehydrogenase I, which is optionally present in overexpressed form, and, where appropriate,
- b) a stop codon chosen from the group consisting of opal, ochre and amber, preferably amber in the rpoS gene, and a t-RNA suppressor chosen from the group
- 10 consisting of opal suppressor, ochre suppressor and amber suppressor, preferably amber suppressor.

Strains which have at least the following features are furthermore suitable in particular for the process according to the invention:

- 15 a) a threonine-insensitive aspartate kinase I - homoserine dehydrogenase I, which is optionally present in overexpressed form,
- b) are not capable, under aerobic culture conditions, of breaking down threonine, preferably by attenuation of
- 20 threonine dehydrogenase,
- c) an at least partial need for isoleucine, and
- d) growth in the presence of at least 5 g/l threonine.

Strains which have at least the following features are very particularly suitable for the process according to the

25 invention:

- a) a threonine-insensitive aspartate kinase I - homoserine dehydrogenase I, which is optionally present in overexpressed form,

- b) are not capable, under aerobic culture conditions, of breaking down threonine, preferably by attenuation of threonine dehydrogenase,
- c) an at least partial need for isoleucine,
- 5 e) growth in the presence of at least 5 g/l threonine, and
- f) a stop codon chosen from the group consisting of opal, ochre and amber, preferably amber in the rpoS gene, and a t-RNA suppressor chosen from the group
- 10 consisting of opal suppressor, ochre suppressor and amber suppressor, preferably amber suppressor.

Moreover, the bacteria employed for the process according to the invention can furthermore have one or more of the following features:

- 15 • Attenuation of phosphoenol pyruvate carboxykinase (PEP carboxykinase), which is coded by the pckA gene, as described, for example, in WO 02/29080,
- Attenuation of phosphoglucose isomerase, which is coded by the pgk gene (Froman et al. Molecular and General
- 20 Genetics 217(1):126-31 (1989)).
- Attenuation of the YtfP gene product, which is coded by the open reading frame ytfP, as described, for example, in WO 02/29080,
- Attenuation of the Yjfa gene product, which is coded by
- 25 the open reading frame yjfa, as described, for example, in WO 02/29080,
- Attenuation of pyruvate oxidase, which is coded by the poxB gene, as described, for example, in WO 02/36797,
- Attenuation of the YjgF gene product, which is coded by
- 30 the open reading frame yjgF, as described, for example,

- in PCT/EP03/14271, The yjgF orf of Escherichia coli has been described by Wasinger VC. and Humphery-Smith I. (FEMS Microbiology Letters 169(2): 375-382 (1998)), Volz K. (Protein Science 8(11): 2428-2437 (1999)) and Parsons
5 et al. (Biochemistry 42(1): 80-89 (2003)). The associated nucleotide or amino acid sequences are available under the Accession Number AE000495 in public databanks. For better clarity, these are shown as SEQ ID NO. 9 and SEQ ID NO. 10.
- 10 • Enhancement of transhydrogenase, which is coded by the genes pntA and pntB, as described, for example, in EP 0 733 712 A1,
- Enhancement of phosphoenol pyruvate synthase, which is coded by the pps gene, as described, for example, in EP 0
15 877 090 A1,
- Enhancement of phosphoenol pyruvate carboxylase, which is coded by the ppc gene, as described, for example, in EP 0 723 011 A1, and
- 20 • Enhancement of the regulator RseB, which is coded by the rseB gene, as described, for example, in EP 1382685. The regulator RseB has been described by Missiakas et al. (Molecular Microbiology 24(2), 355-371 (1997)), De Las Penas et al. (Molecular Microbiology 24(2): 373-385 (1997)) and Collinet et al. (Journal of Biological
25 Chemistry 275(43): 33898-33904 (2000)). The associated nucleotide or amino acid sequences are available under the Accession Number AE000343 in public databanks.
- Enhancement of the galactose proton symporter (= galactose permease), which is coded by the galP gene, as
30 described, for example, in DE 10314618.0. The galP gene and its function have been described by Macpherson et al. (The Journal of Biological Chemistry 258(7): 4390-4396 (1983)) and Venter et al. (The Biochemical Journal 363(Pt

2): 243-252 (2002)). The associated nucleotide or amino acid sequences are available under the Accession Number AE000377 in public databanks.

- Ability to be able to use sucrose as a source of carbon.
5 Genetic determinants for sucrose utilization are described in the prior art, for example in FR-A-2559781, by Debabov (In: Proceedings of the IV International Symposium on Genetics of Industrial Microorganisms 1982. Kodansha Ltd, Tokyo, Japan, p 254-258), Smith and Parsell
10 (Journal of General Microbiology 87,129-140 (1975)) and Livshits et al. (In: Conference on Metabolic Bacterial Plasmids. Tartusk University Press, Tallin, Estonia (1982), p 132-134 and 144-146) and in US 5,705,371. The genetic determinants for sucrose utilization by the
15 strain H155 described by Smith and Parsell were transferred by conjugation into a mutant of Escherichia coli K-12 which is resistant to nalidixic acid and the corresponding transconjugants were deposited on 16th
20 March 2004 at the Deutsche Sammlung für Mikroorganismen und Zellkulturen [German Collection of Microorganisms and Cell Cultures] (Braunschweig, Germany) as DSM 16293. Genetic determinants for sucrose utilization are also contained in the strain 472T23, which is described in US
25 5,631,157 and is obtainable from the ATCC under the name ATCC 9801. A further genetic determinant for sucrose utilization has been described by Bockmann et al. (Molecular and General Genetics 235, 22-32 (1992)) and is known under the name csc system.
- Enhancement of the YedA gene product, which is coded by
30 the open reading frame yedA, as described, for example, in WO 03/044191.
- Growth in the presence of at least 0.1 to 0.5 mM or at least 0.5 to 1 mM borrelidin (borrelidin resistance), as described in US 5,939,307. The borrelidin-resistant

strain kat-13 is obtainable under the Accession Number NRRL B-21593 from the NRRL.

- 5 • Growth in the presence of at least 2 to 2.5 g/l or at least 2.5 to 3 g/l diaminosuccinic acid (diaminosuccinic acid resistance), as described in WO 00/09661. The diaminosuccinic acid-resistant strain DSM 9806 is obtainable under the Accession Number KCCM-10133 from the KCCM.
- 10 • Growth in the presence of at least 30 to 40 mM or at least 40 to 50 mM α -methylserine (α -methylserine resistance), as described in WO 00/09661. The α -methylserine-resistant strain DSM 9806 is obtainable under the Accession Number KCCM-10133 from the KCCM.
- 15 • Growth in the presence of not more than 30 mM or not more than 40 mM or not more than 50 mM fluoropyruvic acid (fluoropyruvic acid sensitivity), as described in WO 00/09661. The fluoropyruvic acid-sensitive strain DSM 9806 is obtainable under the Accession Number KCCM-10133 from the KCCM.
- 20 • Growth in the presence of at least 210 mM or at least 240 mM or at least 270 mM or at least 300 mM L-glutamic acid (glutamic acid resistance), as described in WO 00/09660. The glutamic acid-resistant strain DSM 9807 is obtainable under the Accession Number KCCM-10132 from the KCCM.
- 25 • An at least partial need for methionine. A strain with an at least partial need for methionine is, for example, the strain H-4257, which is described in US 5,017,483 and is obtainable under the Accession Number FERM BP-984 from the National Institute of Advanced Industrial Science and
30 Technology. The need can be compensated by addition of at least 25, 50 or 100 mg/l L-methionine.
- An at least partial need for m-diaminopimelic acid. A strain with an at least partial need for m-diaminopimelic

acid is, for example, the strain H-4257, which is described in US 5,017,483 and is obtainable under the Accession Number FERM BP-984 from the National Institute of Advanced Industrial Science and Technology. The need
5 can be compensated by addition of at least 25, 50 or 100 mg/l m-diaminopimelic acid.

- Growth in the presence of at least 100 mg/l rifampicin (rifampicin resistance), as described in US 4,996,147. The rifampicin-resistant strain H-4581 is obtainable
10 under the Accession Number FERM BP-1411 from the National Institute of Advanced Industrial Science and Technology.
- Growth in the presence of at least 15 g/l L-lysine (lysine resistance), as described in US 4,996,147. The L-lysine-resistant strain H-4581 is obtainable under the
15 Accession Number FERM BP-1411 from the National Institute of Advanced Industrial Science and Technology.
- Growth in the presence of at least 15 g/l methionine (methionine resistance), as described in US 4,996,147. The methionine-resistant strain H-4581 is obtainable
20 under the Accession Number FERM BP-1411 from the National Institute of Advanced Industrial Science and Technology.
- Growth in the presence of at least 15 g/l L-aspartic acid (aspartic acid resistance), as described in US 4,996,147. The aspartic acid-resistant strain H-4581 is obtainable
25 under the Accession Number FERM BP-1411 from the National Institute of Advanced Industrial Science and Technology.
- Enhancement of pyruvate carboxylase, which is coded by the pyc gene. Suitable pyc genes or alleles are, for example, those from *Corynebacterium glutamicum* (WO
30 99/18228, WO 00/39305 and WO 02/31158), *Rhizobium etli* (US 6,455,284) or *Bacillus subtilis* (EP 1092776). If appropriate, the pyc gene from further microorganisms which contain pyruvate carboxylase endogenously, such as,

for example, *Methanobacterium thermoautotrophicum* or *Pseudomonas fluorescens*, can also be used.

If sucrose-containing nutrient media are used, the strains are equipped with genetic determinants for sucrose
5 utilization.

The term "enhancement" in this connection describes the increase in the intracellular activity or concentration of one or more enzymes or proteins in a microorganism which are coded by the corresponding DNA, for example by
10 increasing the number of copies of the open reading frame, gene or allele or open reading frames, genes or alleles by at least one (1) copy, using a potent promoter or a gene or allele which codes for a corresponding enzyme or protein with a high activity, and optionally combining these
15 measures.

For the enhancement measures and also for the attenuation measures, the use of endogenous genes, alleles or open reading frames is in general preferred. "Endogenous genes" or "endogenous nucleotide sequences" are understood as
20 meaning the genes or open reading frames or alleles or nucleotide sequences present in the population of a species.

If plasmids are used to increase the number of copies, these are stabilized, if appropriate, by one or more of the
25 genetic loci chosen from the group consisting of the *parB* locus of the plasmid R1 described by Rasmussen et al. (Molecular and General Genetics 209 (1), 122-128 (1987)), Gerdes et al. (Molecular Microbiology 4 (11), 1807-1818 (1990)) and Thistedt and Gerdes (Journal of Molecular
30 Biology 223 (1), 41-54 (1992)), the *flm* locus of the F plasmid described by Loh et al. (Gene 66 (2), 259-268 (1988)), the *par* locus of the plasmid pSC101 described by Miller et al. (Gene 24 (2-3), 309-315 (1983)), the *cer* locus of the plasmid ColE1 described by Leung et al. (DNA 4 (5),

351-355 (1985), the *par* locus of the plasmid RK2 described by Sobecky et al. (Journal of Bacteriology 178 (7), 2086-2093 (1996)) and Roberts and Helinsky (Journal of Bacteriology 174 (24), 8119-8132 (1992)), the *par* locus of
5 the plasmid RP4 described by Eberl et al. (Molecular Microbiology 12 (1), 131-141 (1994)) and the *parA* locus of the plasmid R1 described by Gerdes and Molin (Journal of Molecular Biology 190 (3), 269- 279 (1986)), Dam and Gerdes (Journal of Molecular Biology 236 (5), 1289- 1298 (1994))
10 and Jensen et al (Proceedings of the National Academy of Sciences USA 95 (15), 8550-8555 (1998)).

By enhancement measures, in particular overexpression, the activity or concentration of the corresponding protein or enzyme is in general increased by at least 10%, 25%, 50%,
15 75%, 100%, 150%, 200%, 300%, 400% or 500%, up to a maximum of 1,000% or 2,000%, based on that of the wild-type protein or the activity or concentration of the protein in the starting microorganism.

To achieve an enhancement, for example, expression of the
20 genes or the catalytic or functional properties of the enzymes or proteins can be increased. The two measures can optionally be combined.

Thus, for example, the number of copies of the corresponding genes can be increased by at least one (1),
25 or the promoter and regulation region or the ribosome binding site upstream of the structural gene can be mutated. Expression cassettes which are incorporated upstream of the structural gene act in the same way. By inducible promoters, it is additionally possible to
30 increase the expression in the course of fermentative L-threonine production. The expression is likewise improved by measures to prolong the life of the m-RNA. Furthermore, the enzyme activity is also increased by preventing the degradation of the enzyme protein. The genes or gene
35 constructs can either be present in plasmids with a varying

number of copies, or can be integrated and amplified in the chromosome. Alternatively, an overexpression of the genes in question can furthermore be achieved by changing the composition of the media and the culture procedure.

- 5 The term "attenuation" in this connection describes the reduction or elimination of the intracellular activity or concentration of one or more enzymes or proteins in a microorganism which are coded by the corresponding DNA, for example by using a weak promoter or an open reading frame
10 or a gene or allele which codes for a corresponding enzyme or protein with a low activity or inactivates the corresponding enzyme or protein or gene and optionally combining these measures.

- By attenuation measures, the activity or concentration of
15 the corresponding protein or enzyme is in general reduced to 0 to 75%, 0 to 50%, 0 to 25%, 0 to 10% or 0 to 5% or 0 to 1% or 0 to 0.1% of the activity or concentration of the wild-type protein or of the activity or concentration of the protein in the starting microorganism.

- 20 To achieve an attenuation, for example, expression of the genes or open reading frames or the catalytic or functional properties of the enzymes or proteins can be reduced or eliminated. The two measures can optionally be combined.

- The reduction in gene expression can take place by suitable
25 culturing, by genetic modification (mutation) of the signal structures of gene expression or also by the antisense-RNA technique. Signal structures of gene expression are, for example, repressor genes, activator genes, operators, promoters, attenuators, ribosome binding sites, the start
30 codon and terminators. The expert can find information in this respect, inter alia, for example, in Jensen and Hammer (Biotechnology and Bioengineering 58: 191-195 (1998)), in Carrier und Keasling (Biotechnology Progress 15: 58-64 (1999)), Franch and Gerdes (Current Opinion in Microbiology

3: 159-164 (2000)) and in known textbooks of genetics and molecular biology, such as, for example, the textbook by Knippers ("Molekulare Genetik [Molecular Genetics]", 6th edition, Georg Thieme Verlag, Stuttgart, Germany, 1995) or
5 that by Winnacker ("Gene und Klone [Genes and Clones]", VCH Verlagsgesellschaft, Weinheim, Germany, 1990).

Mutations which lead to a change or reduction in the catalytic properties of enzyme proteins are known from the prior art. Examples which may be mentioned are the works of
10 Qiu and Goodman (Journal of Biological Chemistry 272: 8611-8617 (1997)), Yano et al. (Proceedings of the National Academy of Sciences of the United States of America 95: 5511-5515 (1998)), Wente and Schachmann (Journal of
Biological Chemistry 266: 20833-20839 (1991)). Summarizing
15 descriptions can be found in known textbooks of genetics and molecular biology, such as e.g. that by Hagemann ("Allgemeine Genetik [General Genetics]", Gustav Fischer Verlag, Stuttgart, 1986).

Possible mutations are transitions, transversions,
20 insertions and deletions of at least one (1) base pair or nucleotide. Depending on the effect of the amino acid exchange caused by the mutation on the enzyme activity, "missense mutations" or "nonsense mutations" are referred to. Missense mutation leads to an exchange of a given amino
25 acid in a protein for another, this being, in particular, a non-conservative amino acid exchange. The functional capacity or activity of the protein is impaired by this means and reduced to a value of 0 to 75%, 0 to 50%, 0 to 25%, 0 to 10%, 0 to 5%, 0 to 1% or 0 to 0.1%. Nonsense
30 mutation leads to a stop codon in the coding region of the gene and therefore to a premature interruption in the translation. Insertions or deletions of at least one base pair in a gene lead to frame shift mutations, which lead to incorrect amino acids being incorporated or translation
35 being interrupted prematurely. If a stop codon is formed in

the coding region as a consequence of the mutation, this also leads to a premature termination of the translation. Deletions of at least one (1) or more codons typically also lead to a complete loss of the enzyme activity or function.

5 Strains which are suitable for the process according to the invention are, inter alia, the strain BKIIM B-3996 described in US 5,175,107, the strain KCCM-10132 described in WO 00/09660, and isoleucine-needing mutants of the strain kat-13 described in WO 98/04715. If appropriate,
10 strains can be adapted to the process according to the invention with the measures mentioned, for example by incorporation of a stop codon into the rpoS gene, for example an amber codon at the site corresponding to position 33 of the amino acid sequence of the RpoS protein,
15 and simultaneous incorporation of a corresponding t-RNA suppressor, for example supE, or by other measures, such as, for example, overexpression of the thrA allele, attenuation of the threonine breakdown which takes place under aerobic culture conditions, introduction of a
20 mutation into the ilvA gene which causes an at least partial need for isoleucine or growth in the presence of at least 5 g/l threonine.

The properties or features mentioned can be transferred into desired strains by transformation, transduction or
25 conjugation.

In the method of transformation, isolated genetic material, typically DNA, is inserted into a recipient strain. In the case of bacteria of the Enterobacteriaceae family, such as e.g. *Escherichia coli*, the DNA for this purpose is
30 incorporated into plasmid or phage DNA in vitro and this is then transferred into the recipient strain. The corresponding methods and working instructions are adequately known in the prior art and are described in detail, for example, in the handbook by J. Sambrook
35 (Molecular Cloning, A Laboratory Manual, 2nd ed., Cold

Spring Harbor Laboratory Press, Cold Spring Harbor, New York, 1989).

Defined mutation can be transferred into suitable strains with the aid of the method of gene or allele exchange using conditionally replicating plasmids. In the case of a defined mutation, at least the position in the chromosome, preferably the exact position of the change in the nucleobase(s) and the nature of the change (substitution, i.e. transition or transversion, insertion or deletion), is known. If appropriate, the corresponding DNA is initially sequenced with the usual methods. A usual method for achieving a gene or allele exchange is that described by Hamilton et al. (Journal of Bacteriology 171: 4617-4622 (1989)), in which the pSC101 derivative pMAK705, which replicates sensitively to heat, is used. Alleles can be transferred from the plasmid into the chromosome with this method. Chromosomal alleles can be transferred to the plasmid in the same manner. Other methods described in the prior art, such as, for example, that of Martinez-Morales et al. (Journal of Bacteriology 181: 7143-7148 (1999)), that of Boyd et al. (Journal of Bacteriology 182: 842-847 (2000)) or the method described in WO 01/77345 can likewise be used.

This method can be employed, inter alia to insert *rpoS* alleles which contain, for example, stop codons, suppressor genes, such as, for example, *supE*, attenuated *tdh* alleles which contain, for example, deletions, attenuated *ilvA* alleles, *thrA* alleles which code for "feed back" resistant aspartate kinase I - homoserine dehydrogenase I variants, the *rhtA23* mutation, attenuated *pck* alleles, attenuated alleles of the *ytfP* ORF, attenuated *yjfA* ORFs, attenuated *poxB* alleles or attenuated *yjgF* ORFs into desired strains.

In the method of transduction, a genetic feature is transferred from a donor strain into a recipient strain using a bacteriophage. This method belongs to the prior art

and is described in textbooks such as, for example, that of E. A. Birge (Bacterial and Bacteriophage Genetics, 4th ed., Springer Verlag, New York, USA, 2000).

In the case of *Escherichia coli*, the bacteriophage P1 is typically used for generalized transduction (Lennox, Virology 1, 190-206 (1955)). A summary of the methods of generalized transduction is given in the article "Generalized Transduction" by M. Masters contained in the textbook by F. C. Neidhard (*Escherichia coli* and *Salmonella* Cellular and Molecular Biology, 2nd ed., ASM Press, Washington, DC, USA, 1996). Practical instructions, for example, are contained in the handbook by J. H. Miller (A Short Course In Bacterial Genetics. A Laboratory Manual and Handbook for *Escherichia coli* and Related Bacteria, Cold Spring Harbor Laboratory Press, New York, USA, 1992) or the handbook by P. Gerhardt "Manual of Methods for General Bacteriology" (American Society for Microbiology, Washington, DC, USA, 1981).

Resistance-imparting or other dominant genetic properties, such as, for example, antibiotics resistance (for example kanamycin resistance, chloramphenicol resistance, rifampicin resistance or borrelidin resistance), resistance to antimetabolites (for example α -amino- β -hydroxyvaleric acid resistance, α -methyl-serine resistance or diaminosuccinic acid resistance), resistance to metabolites (for example threonine resistance, homoserine resistance, glutamic acid resistance, methionine resistance, lysine resistance or aspartic acid resistance) or also the ability to utilize sucrose, can be transferred into suitable recipient strains with the aid of transduction.

The method of transduction is also suitable for inserting so-called non-selectable genetic properties, such as, for example, auxotrophies or needs for amino acids (for example the need for isoleucine, the need for methionine or the need for m-diaminopimelic acid), needs for vitamins or

sensitivity to antimetabolites (for example fluoropyruvic acid sensitivity) into recipient strains. E. coli strains which contain the transposon Tn10 or Tn10kan in an interval of approximately one minute on the chromosome are used for this purpose. These strains are known under the term "Singer Collection" or "Singer/Gross Collection" (Singer et al., Microbiological Reviews 53, 1-24, 1989). These strains are generally available from the E. coli Genetic Stock Center of Yale University (New Haven, CT, USA). Further information is to be found in the article by M. K. B. Berlyn et al. "Linkage Map of Escherichia coli K-12, Edition 9" contained in the textbook by F. C. Neidhard (Escherichia coli and Salmonella Cellular and Molecular Biology, 2nd ed., ASM Press, Washington, DC, USA, 1996). In a similar manner, genetic properties which are not directly selectable (for example fluoropyruvic acid sensitivity, suppressor mutations) and also those whose mutation site is not known, can be transferred into various strains. Instructions in this context are to be found inter alia in the textbook by J. Scaife et al. (Genetics of Bacteria, Academic Press, London, UK, 1985), in the above-mentioned article by M. Masters and in the above-mentioned handbook by J. H. Miller. The tetracycline resistance gene inserted with the transposon Tn10 can optionally be removed again with the method described by Bochner et al. (Journal of Bacteriology 143, 926-933 (1980)).

In the method of conjugation, genetic material is transferred from a donor into a recipient by cell-cell contact. Conjugative transfer of the F factor (F: fertility), conjugative gene transfer using Hfr strains (Hfr: high frequency of recombination) and strains which carry an F' factor (F': F prime) belong to the conventional methods of genetics. Summarizing descriptions are to be found, inter alia, in the standard work by F. C. Neidhard (Escherichia coli and Salmonella Cellular and Molecular Biology, 2nd ed., ASM Press, Washington, DC, USA, 1996).

Practical instructions, for example, are contained in the handbook by J. H. Miller (A Short Course In Bacterial Genetics. A Laboratory Manual and Handbook for Escherichia coli and Related Bacteria, Cold Spring Harbor Laboratory Press, New York, USA, 1992) or the handbook by P. Gerhardt "Manual of Methods for General Bacteriology" (American Society for Microbiology, Washington, DC, USA, 1981). F, F' and Hfr strains are generally available from the E. coli Genetic Stock Center of Yale University (New Haven, CT, USA).

The method of conjugation has been employed, for example, to transfer the mutation thrC1010 described by Thèze and Saint-Girons (Journal of Bacteriology 118, 990-998 (1974)) into the strain MG442 (Debabov, Advances in Biochemical Engineering/Biotechnology 79, 113-136 (2003)). Conjugative plasmids which carry the ability to utilize sucrose are described in the prior art, for example by Schmid et al. (Journal of Bacteriology 151, 68-76 (1982)) or Smith and Parsell (Journal of General Microbiology 87, 129-140 (1975)) and Livshits et al. (In: Conference on Metabolic Bacterial Plasmids. Tartusk University Press, Tallin, Estonia (1982), p 132-134 und 144-146). Thus, Debabov reports (In: Proceedings of the IVth International Symposium on Genetics of Industrial Microorganisms 1982. Kodansha Ltd, Tokyo, Japan, p 254-258) on the construction of threonine-producing strains into which the ability to utilize sucrose has been incorporated with the aid of conjugation.

Patent claims:

1. Process for the preparation of L-threonine using bacteria of the Enterobacteriaceae family which produce L-threonine, characterized in that
 - 5 a) the bacterium is inoculated and cultured in at least a first nutrient medium,
 - b) at least a further nutrient medium or further nutrient media is/are then fed continuously to the culture in one (F1) or several feed streams (F1+),
10 the nutrient medium or the nutrient media comprising at least one source of carbon, at least one source of nitrogen and at least one source of phosphorus, under conditions which allow the formation of L-threonine, and at the same time
15 culture broth is removed from the culture according to the total of the removal stream F2 plus that of the cell-depleted removal stream F3 or according to the total of several removal streams (F2+) plus that of the cell-depleted
20 removal stream F3, which substantially corresponds/correspond to the feed stream (F1) or the total of the feed streams (F1+), wherein
 - c) the cells are completely or partly separated off from the removal stream (F4) with a return flow
25 ratio of $0 < R \leq 1$ by appropriate processes and are recycled into the culturing batch (F5).
2. Process according to claim 1, wherein the culturing step (a) is carried out by the batch process.
3. Process according to claim 1, wherein the culturing
30 step (a) is carried out by the fed batch process, at least one additional nutrient medium being employed.

4. Process according to claim 1, wherein the cell retention is a filtration, centrifugation or sedimentation process.
5. Process according to claim 1, 2, 3 or 4, wherein the L-threonine formed is purified.
6. Process according to claim 1, 2, 3 or 4, wherein in step (b) the biomass is first removed to the extent of at least 90% from the culture removed, and the water is then removed to the extent of at least 90%.
- 10 7. Process according to claim 1 or 2, wherein the further nutrient medium or the further nutrient media is (are) fed in after >0 to 20 hours, with respect to the start of the batch process.
- 15 8. Process according to claim 1 or 3, wherein the further nutrient medium or the further nutrient media is (are) fed in after >0 to 80 hours, with respect to the start of the fed batch process.
- 20 9. Process according to claim 1, wherein the source of carbon is one or more of the compounds chosen from the group consisting of sucrose, molasses from sugar beet or cane sugar, cellulose hydrolysate, arabinose, fructose, glucose, starch hydrolysate, maltose, xylose, acetic acid, ethanol and methanol.
- 25 10. Process according to claim 1, wherein the source of nitrogen is one or more organic nitrogen-containing substances or substance mixtures chosen from the group consisting of peptones, yeast extracts, meat extracts, malt extracts, corn steep liquor, soya bean flour and urea and / or one or more of the inorganic compounds
30 chosen from the group consisting of ammonia, ammonium-containing salts and salts of nitric acid.

11. Process according to claim 10, wherein the ammonium-containing salts and salts of nitric acid are ammonium sulfate, ammonium chloride, ammonium phosphate, ammonium carbonate, ammonium nitrate, potassium nitrate and potassium sodium nitrate.
5
12. Process according to claim 1, wherein the source of phosphorus is alkali metal or alkaline earth metal salts of phosphoric acid or polymers thereof or of phytic acid.
- 10 13. Process according to claim 12, wherein the alkali metal salts of phosphoric acid are potassium dihydrogen phosphate or dipotassium hydrogen phosphate or the corresponding sodium-containing salts.
14. Process according to claim 1, wherein the speed of the removal stream F2 or the removal streams F2+ corresponds to 80% - 120%, 90% - 110% of the feed stream F1 or of the total of the feed streams F1+.
15
15. Process according to claim 1, wherein the bacteria of the Enterobacteriaceae family are the species Escherichia coli.
20
16. Process according to claim 1, wherein the bacterium of the Enterobacteriaceae family contains at least one thrA gene or allele which codes for a threonine-insensitive aspartate kinase I - homoserine dehydrogenase I.
25
17. Process according to claim 1, wherein the bacterium of the Enterobacteriaceae family contains a stop codon chosen from the group consisting of opal, ochre and amber, preferably amber, in the rpoS gene and a t-RNA suppressor chosen from the group consisting of opal suppressor, ochre suppressor and amber suppressor, preferably amber suppressor.
30

18. Process according to claim 1, wherein the feed stream (F1) or the total of the feed streams (F1+) is fed in at a rate corresponding to an average residence time of less than 30 hours, less than 25, less than 20 hours.
19. Process according to claim 1, wherein in the nutrient medium fed in or the nutrient media fed in a phosphorus to carbon ratio (P/C ratio) of not more than 4; of not more than 3; of not more than 2; of not more than 1.5; of not more than 1; of not more than 0.7; of not more than 0.5; of not more than 0.48; of not more than 0.46; of not more than 0.44; of not more than 0.42; of not more than 0.40; of not more than 0.38; of not more than 0.36; of not more than 0.34; of not more than 0.32; or of not more than 0.30 is established.
20. Process according to claim 1, wherein the culture broth removed is provided with oxygen or an oxygen-containing gas until the concentration of the source of carbon falls below 2 g/l; below 1 g/l; below 0.5 g/l.
21. Process according to claim 20, wherein the L-threonine formed is purified.
22. Process according to claim 20, wherein in step (b) the biomass is first removed to the extent of at least 90% from the culture removed, and the water is then removed.
23. Process according to claim 1, 2, 3 or 4, wherein the concentration of the source of carbon during the culture is adjusted to not more than 30, 20, 10 or 5 g/l.

24. Process according to claim 1, 2, 3 or 4, wherein the concentration of the source of carbon during the culture is adjusted to not more than 10 g/l.
- 5 25. Process according to claim 1, 2, 3 or 4, wherein the concentration of the source of carbon during the culture is adjusted to not more than 5 g/l.
26. Process according to claim 1, 2, 3 or 4, wherein the concentration of the source of carbon during the culture is adjusted to not more than 2 g/l.
- 10 27. Process according to claim 1, 2, 3 or 4, wherein the yield of L-threonine formed, based on the source of carbon employed, is at least 31%.
28. Process according to claim 1, 2, 3 or 4, wherein the yield of L-threonine formed, based on the source of carbon employed, is at least 37%.
- 15 29. Process according to claim 1, 2, 3 or 4, wherein the yield of L-threonine formed, based on the source of carbon employed, is at least 42%.
30. Process according to claim 1, 2, 3 or 4, wherein the yield of L-threonine formed, based on the source of carbon employed, is at least 48%.
- 20 31. Process according to claim 1, 2, 3 or 4, wherein L-threonine is formed with a space/time yield of 5.0 to more than 8.0 g/l per h.
- 25 32. Process according to claim 1, 2, 3 or 4, wherein L-threonine is formed with a space/time yield of 3.5 to more than 5.0 g/l per h.
33. Process according to claim 1, 2, 3 or 4, wherein L-threonine is formed with a space/time yield of 2.5 to more than 3.5 g/l per h.
- 30

34. Process according to claim 1, wherein a fed batch process is used in the culturing step (a), and in that L-threonine is formed with a space/time yield of at least 2.5 to 5.0 g/l per h.
- 5 35. Sucrose-utilizing transconjugants of *Escherichia coli* K-12 deposited as DSM 16293 at the Deutsche Sammlung für Mikroorganismen und Zellkulturen [German Collection of Microorganisms and Cell Cultures] (Braunschweig, Germany).
- 10 36. Process according to claim 1, 2 or 3, wherein strains which have at least the following features are employed:
- 15 a) a threonine-insensitive aspartate kinase I - homoserine dehydrogenase I, which is optionally present in overexpressed form, and, where appropriate,
- 20 b) a stop codon chosen from the group consisting of opal, ochre and amber, preferably amber in the *rpoS* gene, and a t-RNA suppressor chosen from the group consisting of opal suppressor, ochre suppressor and amber suppressor.
37. Process according to claim 1, 2 or 3, wherein strains which have at least the following features are employed:
- 25 a) a threonine-insensitive aspartate kinase I - homoserine dehydrogenase I, which is optionally present in overexpressed form,
- b) are not capable, under aerobic culture conditions, of breaking down threonine,
- 30 c) an at least partial need for isoleucine, and

d) growth in the presence of at least 5 g/l threonine.

38. Process according to claim 1, 2 or 3, wherein strains which have at least the following features are employed:

- a) a threonine-insensitive aspartate kinase I - homoserine dehydrogenase I, which is optionally present in overexpressed form,
- b) a stop codon chosen from the group consisting of opal, ochre and amber, preferably amber in the rpoS gene, and a t-RNA suppressor chosen from the group consisting of opal suppressor, ochre suppressor and amber suppressor,
- c) are not capable, under aerobic culture conditions, of breaking down threonine, preferably by attenuation of threonine dehydrogenase,
- d) an at least partial need for isoleucine, and
- e) growth in the presence of at least 5 g/l threonine.

39. Process according to claim 36, 37 or 38, wherein the strain employed additionally contains one or more of the features chosen from the group consisting of


- 39.1 attenuation of phosphoenol pyruvate carboxykinase, which is coded by the pckA gene,
- 39.2 attenuation of phosphoglucose isomerase, which is coded by the pgi gene,
- 39.3 attenuation of the YtfP gene product, which is coded by the open reading frame ytfP,

- 39.4 attenuation of the YjfA gene product, which is coded by the open reading frame yjfA,
- 39.5 attenuation of pyruvate oxidase, which is coded by the poxB gene,
- 5 39.6 attenuation of the YjgF gene product, which is coded by the open reading frame yjgF,
- 39.7 enhancement of transhydrogenase, which is coded by the genes pntA and pntB,
- 39.8 enhancement of phosphoenol pyruvate synthase, which is coded by the pps gene,
- 10 39.9 enhancement of phosphoenol pyruvate carboxylase, which is coded by the ppc gene.
- 39.10 enhancement of the regulator RseB, which is coded by the rseB gene,
- 15 39.11 enhancement of the galactose proton symporter, which is coded by the galP gene,
- 39.12 ability to be able to use sucrose as a source of carbon,
- 39.13 enhancement of the YedA gene product, which is coded by the open reading frame yedA,
- 20 39.14 growth in the presence of at least 0.1 to 0.5 mM or at least 0.5 to 1 mM borrelidin (borrelidin resistance),
- 39.15 growth in the presence of at least 2 to 2.5 g/l or at least 2.5 to 3 g/l diaminosuccinic acid (diaminosuccinic acid resistance),
- 25 39.16 growth in the presence of at least 30 to 40 mM or at least 40 to 50 mM α -methylserine (α -methylserine resistance),

- 39.17 growth in the presence of not more than 30 mM or not more than 40 mM or not more than 50 mM fluoropyruvic acid (fluoropyruvic acid sensitivity),
- 5 39.18 growth in the presence of at least 210 mM or at least 240 mM or at least 270 mM or at least 300 mM L-glutamic acid (glutamic acid resistance),
- 39.19 an at least partial need for methionine,
- 10 39.20 an at least partial need for m-diaminopimelic acid,
- 39.21 growth in the presence of at least 100 mg/l rifampicin (rifampicin resistance),
- 15 39.22 growth in the presence of at least 15 g/l L-lysine (lysine resistance),
- 39.23 growth in the presence of at least 15 g/l methionine (methionine resistance),
- 39.24 growth in the presence of at least 15 g/l L-aspartic acid (aspartic acid resistance), and
- 20 39.25 enhancement of pyruvate carboxylase, which is coded by the pyc gene.

1/1

Figure 1



SEQUENCE LISTING

5 <110> Degussa AG

 <120> Process for the preparation of L-threonine

10 <130> 030219BT

 <160> 10

15 <170> PatentIn version 3.1

20 <210> 1
 <211> 993
 <212> DNA
 <213> Escherichia coli

25 <220>
 <221> CDS
 <222> (1)..(990)
 <223> rpoS gene

30 <400> 1

35 atg agt cag aat acg ctg aaa gtt cat gat tta aat gaa gat gcg gaa 48
 Met Ser Gln Asn Thr Leu Lys Val His Asp Leu Asn Glu Asp Ala Glu
 1 5 10 15

40 ttt gat gag aac gga gtt gag gtt ttt gac gaa aag gcc tta gta gaa 96
 Phe Asp Glu Asn Gly Val Glu Val Phe Asp Glu Lys Ala Leu Val Glu
 20 25 30

45 cag gaa ccc agt gat aac gat ttg gcc gaa gag gaa ctg tta tcg cag 144
 Gln Glu Pro Ser Asp Asn Asp Leu Ala Glu Glu Glu Leu Leu Ser Gln
 35 40 45

50 gga gcc aca cag cgt gtg ttg gac gcg act cag ctt tac ctt ggt gag 192
 Gly Ala Thr Gln Arg Val Leu Asp Ala Thr Gln Leu Tyr Leu Gly Glu
 50 55 60

55 att ggt tat tca cca ctg tta acg gcc gaa gaa gaa gtt tat ttt gcg 240
 Ile Gly Tyr Ser Pro Leu Leu Thr Ala Glu Glu Glu Val Tyr Phe Ala
 65 70 75 80

60 cgt cgc gca ctg cgt gga gat gtc gcc tct cgc cgc cgg atg atc gag 288
 Arg Arg Ala Leu Arg Gly Asp Val Ala Ser Arg Arg Arg Met Ile Glu
 85 90 95

60 agt aac ttg cgt ctg gtg gta aaa att gcc cgc cgt tat ggc aat cgt 336
 Ser Asn Leu Arg Leu Val Val Lys Ile Ala Arg Arg Tyr Gly Asn Arg
 100 105 110

2

	ggt ctg gcg ttg ctg gac ctt atc gaa gag ggc aac ctg ggg ctg atc	384
	Gly Leu Ala Leu Leu Asp Leu Ile Glu Glu Gly Asn Leu Gly Leu Ile	
	115 120 125	
5	cgc gcg gta gag aag ttt gac ccg gaa cgt ggt ttc cgc ttc tca aca	432
	Arg Ala Val Glu Lys Phe Asp Pro Glu Arg Gly Phe Arg Phe Ser Thr	
	130 135 140	
10	tac gca acc tgg tgg att cgc cag acg att gaa cgg gcg att atg aac	480
	Tyr Ala Thr Trp Trp Ile Arg Gln Thr Ile Glu Arg Ala Ile Met Asn	
	145 150 155 160	
15	caa acc cgt act att cgt ttg ccg att cac atc gta aag gag ctg aac	528
	Gln Thr Arg Thr Ile Arg Leu Pro Ile His Ile Val Lys Glu Leu Asn	
	165 170 175	
20	gtt tac ctg cga acc gca cgt gag ttg tcc cat aag ctg gac cat gaa	576
	Val Tyr Leu Arg Thr Ala Arg Glu Leu Ser His Lys Leu Asp His Glu	
	180 185 190	
25	cca agt gcg gaa gag atc gca gag caa ctg gat aag cca gtt gat gac	624
	Pro Ser Ala Glu Glu Ile Ala Glu Gln Leu Asp Lys Pro Val Asp Asp	
	195 200 205	
30	gtc agc cgt atg ctt cgt ctt aac gag cgc att acc tcg gta gac acc	672
	Val Ser Arg Met Leu Arg Leu Asn Glu Arg Ile Thr Ser Val Asp Thr	
	210 215 220	
35	ccg ctg ggt ggt gat tcc gaa aaa gcg ttg ctg gac atc ctg gcc gat	720
	Pro Leu Gly Gly Asp Ser Glu Lys Ala Leu Leu Asp Ile Leu Ala Asp	
	225 230 235 240	
40	gaa aaa gag aac ggt ccg gaa gat acc acg caa gat gac gat atg aag	768
	Glu Lys Glu Asn Gly Pro Glu Asp Thr Thr Gln Asp Asp Asp Met Lys	
	245 250 255	
45	cag agc atc gtc aaa tgg ctg ttc gag ctg aac gcc aaa cag cgt gaa	816
	Gln Ser Ile Val Lys Trp Leu Phe Glu Leu Asn Ala Lys Gln Arg Glu	
	260 265 270	
50	gtg ctg gca cgt cga ttc ggt ttg ctg ggg tac gaa gcg gca aca ctg	864
	Val Leu Ala Arg Arg Phe Gly Leu Leu Gly Tyr Glu Ala Ala Thr Leu	
	275 280 285	
55	gaa gat gta ggt cgt gaa att ggc ctc acc cgt gaa cgt gtt cgc cag	912
	Glu Asp Val Gly Arg Glu Ile Gly Leu Thr Arg Glu Arg Val Arg Gln	
	290 295 300	
60	att cag gtt gaa ggc ctg cgc cgt ttg cgc gaa atc ctg caa acg cag	960
	Ile Gln Val Glu Gly Leu Arg Arg Leu Arg Glu Ile Leu Gln Thr Gln	
	305 310 315 320	
65	ggg ctg aat atc gaa gcg ctg ttc cgc gag taa	993
	Gly Leu Asn Ile Glu Ala Leu Phe Arg Glu	
	325 330	
	<210> 2	
	<211> 330	
	<212> PRT	
	<213> Escherichia coli	

```

<400> 2
Met Ser Gln Asn Thr Leu Lys Val His Asp Leu Asn Glu Asp Ala Glu
1      5      10      15

5  Phe Asp Glu Asn Gly Val Glu Val Phe Asp Glu Lys Ala Leu Val Glu
      20      25      30

      Gln Glu Pro Ser Asp Asn Asp Leu Ala Glu Glu Glu Leu Leu Ser Gln
      35      40      45

10  Gly Ala Thr Gln Arg Val Leu Asp Ala Thr Gln Leu Tyr Leu Gly Glu
      50      55      60

      Ile Gly Tyr Ser Pro Leu Leu Thr Ala Glu Glu Glu Val Tyr Phe Ala
15  65      70      75      80

      Arg Arg Ala Leu Arg Gly Asp Val Ala Ser Arg Arg Arg Met Ile Glu
      85      90      95

20  Ser Asn Leu Arg Leu Val Val Lys Ile Ala Arg Arg Tyr Gly Asn Arg
      100      105      110

      Gly Leu Ala Leu Leu Asp Leu Ile Glu Glu Gly Asn Leu Gly Leu Ile
      115      120      125

25  Arg Ala Val Glu Lys Phe Asp Pro Glu Arg Gly Phe Arg Phe Ser Thr
      130      135      140

      Tyr Ala Thr Trp Trp Ile Arg Gln Thr Ile Glu Arg Ala Ile Met Asn
30  145      150      155      160

      Gln Thr Arg Thr Ile Arg Leu Pro Ile His Ile Val Lys Glu Leu Asn
      165      170      175

35  Val Tyr Leu Arg Thr Ala Arg Glu Leu Ser His Lys Leu Asp His Glu
      180      185      190

      Pro Ser Ala Glu Glu Ile Ala Glu Gln Leu Asp Lys Pro Val Asp Asp
      195      200      205

40  Val Ser Arg Met Leu Arg Leu Asn Glu Arg Ile Thr Ser Val Asp Thr
      210      215      220

      Pro Leu Gly Gly Asp Ser Glu Lys Ala Leu Leu Asp Ile Leu Ala Asp
45  225      230      235      240

      Glu Lys Glu Asn Gly Pro Glu Asp Thr Thr Gln Asp Asp Asp Met Lys
      245      250      255

50  Gln Ser Ile Val Lys Trp Leu Phe Glu Leu Asn Ala Lys Gln Arg Glu
      260      265      270

      Val Leu Ala Arg Arg Phe Gly Leu Leu Gly Tyr Glu Ala Ala Thr Leu
      275      280      285

55  Glu Asp Val Gly Arg Glu Ile Gly Leu Thr Arg Glu Arg Val Arg Gln
      290      295      300

      Ile Gln Val Glu Gly Leu Arg Arg Leu Arg Glu Ile Leu Gln Thr Gln
60  305      310      315      320

```


Gly Leu Asn Ile Glu Ala Leu Phe Arg Glu
 325 330

5 <210> 3
 <211> 993
 <212> DNA
 <213> Escherichia coli

10 <220>
 <221> Allele
 <222> (1)..(990)
 <223> rpoS allele

15 <220>
 <221> misc_feature
 <222> (97)..(99)
 <223> amber codon

20 <400> 3
 atgagtcaga atacgctgaa agttcatgat ttaaataaag atgcggaatt tgatgagaac 60

25 ggagttgagg tttttgacga aaaggcctta gtagaatagg aaccacagtga taacgatttg 120
 gccgaagagg aactgttata gcaggagacc acacagcgtg tggtggacgc gactcagctt 180

30 taccttggtg agattggtta ttcaccactg ttaacggccg aagaagaagt ttattttgcg 240
 cgtcgcgcac tgcgtggaga tgcgcctct cgccgccgga tgatcgagag taacttgcgt 300
 ctggtggttaa aaattgcccg ccgttatggc aatcgtggtc tggcgttgct ggaccttata 360

35 gaagagggca acctggggct gatccgcgcg gtagagaagt ttgaccggga acgtgggtttc 420
 cgcttctcaa catacgcaac ctggtggatt cgccagacga ttgaacgggc gattatgaac 480

40 caaaccgcgt ctattcggtt gccgattcac atcgtaaagg agctgaacgt ttacctgcga 540
 accgcacgtg agttgtccca taagctggac catgaaccaa gtgcggaaga gatcgagag 600
 caactggata agccagttga tgacgtcagc cgtatgcttc gtcttaacga ggcattacc 660

45 tcggtagaca ccccgctggg tgggtattcc gaaaaagcgt tgctggacat cctggccgat 720
 gaaaaagaga acggtccgga agataccacg caagatgacg atatgaagca gagcatcgtc 780

50 aaatggctgt tcgagctgaa cgccaaacag cgtgaagtgc tggcacgtcg attcggtttg 840
 ctgggggtacg aagcggcaac actggaagat gtaggtcgtg aaattggcct caccgtgaa 900
 cgtgttcgcc agattcaggt tgaaggcctg cgccgtttgc gcgaaatcct gcaaacgcag 960

55 gggctgaata tcgaagcgt gttccgcgag taa 993

60 <210> 4
 <211> 75
 <212> DNA
 <213> Escherichia coli

5

<220>
 <221> tRNA
 <222> (1)..(75)
 <223> supE allele
 5

 <400> 4
 tgggggtatcgc ccaagcggta aggcaccgga ttctaattcc ggcattccga gggttcgaatc 60
 10 ctcgtacccc agcca 75

 <210> 5
 <211> 1545
 <212> DNA
 15 <213> Escherichia coli

 <220>
 <221> CDS
 20 <222> (1)..(1542)
 <223> ilvA gene

 <400> 5
 25 atg gct gac tcg caa ccc ctg tcc ggt gct ccg gaa ggt gcc gaa tat 48
 Met Ala Asp Ser Gln Pro Leu Ser Gly Ala Pro Glu Gly Ala Glu Tyr
 1 5 10 15

 tta aga gca gtg ctg cgc gcg ccg gtt tac gag gcg gcg cag gtt acg 96
 30 Leu Arg Ala Val Leu Arg Ala Pro Val Tyr Glu Ala Ala Gln Val Thr
 20 25 30

 ccg cta caa aaa atg gaa aaa ctg tcg tcg cgt ctt gat aac gtc att 144
 Pro Leu Gln Lys Met Glu Lys Leu Ser Ser Arg Leu Asp Asn Val Ile
 35 35 40 45

 ctg gtg aag cgc gaa gat cgc cag cca gtg cac agc ttt aag ctg cgc 192
 Leu Val Lys Arg Glu Asp Arg Gln Pro Val His Ser Phe Lys Leu Arg
 50 55 60

 40 ggc gca tac gcc atg atg gcg ggc ctg acg gaa gaa cag aaa gcg cac 240
 Gly Ala Tyr Ala Met Met Ala Gly Leu Thr Glu Glu Gln Lys Ala His
 65 70 75 80

 ggc gtg atc act gct tct gcg ggt aac cac gcg cag ggc gtc gcg ttt 288
 45 Gly Val Ile Thr Ala Ser Ala Gly Asn His Ala Gln Gly Val Ala Phe
 85 90 95

 tct tct gcg cgg tta ggc gtg aag gcc ctg atc gtt atg cca acc gcc 336
 50 Ser Ser Ala Arg Leu Gly Val Lys Ala Leu Ile Val Met Pro Thr Ala
 100 105 110

 acc gcc gac atc aaa gtc gac gcg gtg cgc ggc ttc ggc ggc gaa gtg 384
 Thr Ala Asp Ile Lys Val Asp Ala Val Arg Gly Phe Gly Gly Glu Val
 115 120 125
 55
 ctg ctc cac ggc gcg aac ttt gat gaa gcg aaa gcc aaa gcg atc gaa 432
 Leu Leu His Gly Ala Asn Phe Asp Glu Ala Lys Ala Lys Ala Ile Glu
 130 135 140

 60 ctg tca cag cag cag ggg ttc acc tgg gtg ccg ccg ttc gac cat ccg 480
 Leu Ser Gln Gln Gln Gly Phe Thr Trp Val Pro Phe Asp His Pro
 145 150 155 160

	atg gtg att gcc ggg caa ggc acg ctg gcg ctg gaa ctg ctc cag cag	528
	Met Val Ile Ala Gly Gln Gly Thr Leu Ala Leu Glu Leu Leu Gln Gln	
5	165 170 175	
	gac gcc cat ctc gac cgc gta ttt gtg cca gtc ggc ggc ggc ggt ctg	576
	Asp Ala His Leu Asp Arg Val Phe Val Pro Val Gly Gly Gly Gly Leu	
	180 185 190	
10	gct gct ggc gtg gcg gtg ctg atc aaa caa ctg atg ccg caa atc aaa	624
	Ala Ala Gly Val Ala Val Leu Ile Lys Gln Leu Met Pro Gln Ile Lys	
	195 200 205	
15	gtg atc gcc gta gaa gcg gaa gac tcc gcc tgc ctg aaa gca gcg ctg	672
	Val Ile Ala Val Glu Ala Glu Asp Ser Ala Cys Leu Lys Ala Ala Leu	
	210 215 220	
20	gat gcg ggt cat ccg gtt gat ctg ccg cgc gta ggg cta ttt gct gaa	720
	Asp Ala Gly His Pro Val Asp Leu Pro Arg Val Gly Leu Phe Ala Glu	
	225 230 235 240	
25	ggc gta gcg gta aaa cgc atc ggt gac gaa acc ttc cgt tta tgc cag	768
	Gly Val Ala Val Lys Arg Ile Gly Asp Glu Thr Phe Arg Leu Cys Gln	
	245 250 255	
	gag tat ctc gac gac atc atc acc gtc gat agc gat gcg atc tgt gcg	816
	Glu Tyr Leu Asp Asp Ile Ile Thr Val Asp Ser Asp Ala Ile Cys Ala	
	260 265 270	
30	gcg atg aag gat tta ttc gaa gat gtg cgc gcg gtg gcg gaa ccc tct	864
	Ala Met Lys Asp Leu Phe Glu Asp Val Arg Ala Val Ala Glu Pro Ser	
	275 280 285	
35	ggc gcg ctg gcg ctg gcg gga atg aaa aaa tat atc gcc ctg cac aac	912
	Gly Ala Leu Ala Leu Ala Gly Met Lys Lys Tyr Ile Ala Leu His Asn	
	290 295 300	
40	att cgc ggc gaa cgg ctg gcg cat att ctt tcc ggt gcc aac gtg aac	960
	Ile Arg Gly Glu Arg Leu Ala His Ile Leu Ser Gly Ala Asn Val Asn	
	305 310 315 320	
45	ttc cac ggc ctg cgc tac gtc tca gaa cgc tgc gaa ctg ggc gaa cag	1008
	Phe His Gly Leu Arg Tyr Val Ser Glu Arg Cys Glu Leu Gly Glu Gln	
	325 330 335	
	cgt gaa gcg ttg ttg gcg gtg acc att ccg gaa gaa aaa ggc agc ttc	1056
	Arg Glu Ala Leu Leu Ala Val Thr Ile Pro Glu Glu Lys Gly Ser Phe	
	340 345 350	
50	ctc aaa ttc tgc caa ctg ctt ggc ggc cgt tcg gtc acc gag ttc aac	1104
	Leu Lys Phe Cys Gln Leu Leu Gly Gly Arg Ser Val Thr Glu Phe Asn	
	355 360 365	
55	tac cgt ttt gcc gat gcc aaa aac gcc tgc atc ttt gtc ggt gtg cgc	1152
	Tyr Arg Phe Ala Asp Ala Lys Asn Ala Cys Ile Phe Val Gly Val Arg	
	370 375 380	
60	ctg agc cgc ggc ctc gaa gag cgc aaa gaa att ttg cag atg ctc aac	1200
	Leu Ser Arg Gly Leu Glu Glu Arg Lys Glu Ile Leu Gln Met Leu Asn	
	385 390 395 400	

7

	gac ggc ggc tac agc gtg gtt gat ctc tcc gac gac gaa atg gcg aag	1248
	Asp Gly Gly Tyr Ser Val Val Asp Leu Ser Asp Asp Glu Met Ala Lys	
	405 410 415	
5	cta cac gtg cgc tat atg gtc ggc gga cgt cca tcg cat ccg ttg cag	1296
	Leu His Val Arg Tyr Met Val Gly Gly Arg Pro Ser His Pro Leu Gln	
	420 425 430	
10	gaa cgc ctc tac agc ttc gaa ttc ccg gaa tca ccg ggc gcg ctg ctg	1344
	Glu Arg Leu Tyr Ser Phe Glu Phe Pro Glu Ser Pro Gly Ala Leu Leu	
	435 440 445	
15	cgc ttc ctc aac acg ctg ggt acg tac tgg aac att tct ttg ttc cac	1392
	Arg Phe Leu Asn Thr Leu Gly Thr Tyr Trp Asn Ile Ser Leu Phe His	
	450 455 460	
20	tat cgc agc cat ggc acc gac tac ggg cgc gta ctg gcg gcg ttc gaa	1440
	Tyr Arg Ser His Gly Thr Asp Tyr Gly Arg Val Leu Ala Ala Phe Glu	
	465 470 475 480	
	ctt ggc gac cat gaa ccg gat ttc gaa acc cgg ctg aat gag ctg ggc	1488
	Leu Gly Asp His Glu Pro Asp Phe Glu Thr Arg Leu Asn Glu Leu Gly	
	485 490 495	
25	tac gat tgc cac gac gaa acc aat aac ccg gcg ttc agg ttc ttt ttg	1536
	Tyr Asp Cys His Asp Glu Thr Asn Asn Pro Ala Phe Arg Phe Phe Leu	
	500 505 510	
30	gcg ggt tag	1545
	Ala Gly	
35	<210> 6	
	<211> 514	
	<212> PRT	
	<213> Escherichia coli	
40	<400> 6	
	Met Ala Asp Ser Gln Pro Leu Ser Gly Ala Pro Glu Gly Ala Glu Tyr	
	1 5 10 15	
45	Leu Arg Ala Val Leu Arg Ala Pro Val Tyr Glu Ala Ala Gln Val Thr	
	20 25 30	
	Pro Leu Gln Lys Met Glu Lys Leu Ser Ser Arg Leu Asp Asn Val Ile	
	35 40 45	
50	Leu Val Lys Arg Glu Asp Arg Gln Pro Val His Ser Phe Lys Leu Arg	
	50 55 60	
	Gly Ala Tyr Ala Met Met Ala Gly Leu Thr Glu Glu Gln Lys Ala His	
	65 70 75 80	
55	Gly Val Ile Thr Ala Ser Ala Gly Asn His Ala Gln Gly Val Ala Phe	
	85 90 95	
60	Ser Ser Ala Arg Leu Gly Val Lys Ala Leu Ile Val Met Pro Thr Ala	
	100 105 110	
	Thr Ala Asp Ile Lys Val Asp Ala Val Arg Gly Phe Gly Gly Glu Val	
	115 120 125	

	Leu	Leu	His	Gly	Ala	Asn	Phe	Asp	Glu	Ala	Lys	Ala	Lys	Ala	Ile	Glu
	130						135					140				
5	Leu	Ser	Gln	Gln	Gln	Gly	Phe	Thr	Trp	Val	Pro	Pro	Phe	Asp	His	Pro
	145					150					155					160
	Met	Val	Ile	Ala	Gly	Gln	Gly	Thr	Leu	Ala	Leu	Glu	Leu	Leu	Gln	Gln
					165					170					175	
10	Asp	Ala	His	Leu	Asp	Arg	Val	Phe	Val	Pro	Val	Gly	Gly	Gly	Gly	Leu
				180					185					190		
	Ala	Ala	Gly	Val	Ala	Val	Leu	Ile	Lys	Gln	Leu	Met	Pro	Gln	Ile	Lys
15			195					200					205			
	Val	Ile	Ala	Val	Glu	Ala	Glu	Asp	Ser	Ala	Cys	Leu	Lys	Ala	Ala	Leu
	210						215					220				
20	Asp	Ala	Gly	His	Pro	Val	Asp	Leu	Pro	Arg	Val	Gly	Leu	Phe	Ala	Glu
	225					230					235					240
	Gly	Val	Ala	Val	Lys	Arg	Ile	Gly	Asp	Glu	Thr	Phe	Arg	Leu	Cys	Gln
					245					250					255	
25	Glu	Tyr	Leu	Asp	Asp	Ile	Ile	Thr	Val	Asp	Ser	Asp	Ala	Ile	Cys	Ala
				260					265					270		
	Ala	Met	Lys	Asp	Leu	Phe	Glu	Asp	Val	Arg	Ala	Val	Ala	Glu	Pro	Ser
30			275					280					285			
	Gly	Ala	Leu	Ala	Leu	Ala	Gly	Met	Lys	Lys	Tyr	Ile	Ala	Leu	His	Asn
	290						295					300				
35	Ile	Arg	Gly	Glu	Arg	Leu	Ala	His	Ile	Leu	Ser	Gly	Ala	Asn	Val	Asn
	305					310					315					320
	Phe	His	Gly	Leu	Arg	Tyr	Val	Ser	Glu	Arg	Cys	Glu	Leu	Gly	Glu	Gln
					325					330					335	
40	Arg	Glu	Ala	Leu	Leu	Ala	Val	Thr	Ile	Pro	Glu	Glu	Lys	Gly	Ser	Phe
				340					345					350		
	Leu	Lys	Phe	Cys	Gln	Leu	Leu	Gly	Gly	Arg	Ser	Val	Thr	Glu	Phe	Asn
45			355					360					365			
	Tyr	Arg	Phe	Ala	Asp	Ala	Lys	Asn	Ala	Cys	Ile	Phe	Val	Gly	Val	Arg
	370						375					380				
50	Leu	Ser	Arg	Gly	Leu	Glu	Glu	Arg	Lys	Glu	Ile	Leu	Gln	Met	Leu	Asn
	385					390					395					400
	Asp	Gly	Gly	Tyr	Ser	Val	Val	Asp	Leu	Ser	Asp	Asp	Glu	Met	Ala	Lys
					405					410					415	
55	Leu	His	Val	Arg	Tyr	Met	Val	Gly	Gly	Arg	Pro	Ser	His	Pro	Leu	Gln
				420					425					430		
	Glu	Arg	Leu	Tyr	Ser	Phe	Glu	Phe	Pro	Glu	Ser	Pro	Gly	Ala	Leu	Leu
60			435					440					445			

Arg Phe Leu Asn Thr Leu Gly Thr Tyr Trp Asn Ile Ser Leu Phe His
 450 455 460
 5 Tyr Arg Ser His Gly Thr Asp Tyr Gly Arg Val Leu Ala Ala Phe Glu
 465 470 475 480
 Leu Gly Asp His Glu Pro Asp Phe Glu Thr Arg Leu Asn Glu Leu Gly
 485 490 495
 10 Tyr Asp Cys His Asp Glu Thr Asn Asn Pro Ala Phe Arg Phe Phe Leu
 500 505 510
 Ala Gly
 15
 <210> 7
 <211> 1545
 <212> DNA
 <213> Escherichia coli
 20
 <220>
 <221> CDS
 <222> (1)..(1542)
 25 <223> ilvA allele
 <220>
 <221> mutation
 30 <222> (856)..(856)
 <223>
 <400> 7
 35 atg gct gac tcg caa ccc ctg tcc ggt gct ccg gaa ggt gcc gaa tat 48
 Met Ala Asp Ser Gln Pro Leu Ser Gly Ala Pro Glu Gly Ala Glu Tyr
 1 5 10 15
 40 tta aga gca gtg ctg cgc gcg ccg gtt tac gag gcg gcg cag gtt acg 96
 Leu Arg Ala Val Leu Arg Ala Pro Val Tyr Glu Ala Ala Gln Val Thr
 20 25 30
 ccg cta caa aaa atg gaa aaa ctg tcg tcg cgt ctt gat aac gtc att 144
 Pro Leu Gln Lys Met Glu Lys Leu Ser Ser Arg Leu Asp Asn Val Ile
 35 40 45
 45 ctg gtg aag cgc gaa gat cgc cag cca gtg cac agc ttt aag ctg cgc 192
 Leu Val Lys Arg Glu Asp Arg Gln Pro Val His Ser Phe Lys Leu Arg
 50 55 60
 50 ggc gca tac gcc atg atg gcg ggc ctg acg gaa gaa cag aaa gcg cac 240
 Gly Ala Tyr Ala Met Met Ala Gly Leu Thr Glu Glu Gln Lys Ala His
 65 70 75 80
 55 ggc gtg atc act gct tct gcg ggt aac cac gcg cag ggc gtc gcg ttt 288
 Gly Val Ile Thr Ala Ser Ala Gly Asn His Ala Gln Gly Val Ala Phe
 85 90 95
 60 tct tct gcg cgg tta ggc gtg aag gcc ctg atc gtt atg cca acc gcc 336
 Ser Ser Ala Arg Leu Gly Val Lys Ala Leu Ile Val Met Pro Thr Ala
 100 105 110

10

	acc gcc gac atc aaa gtc gac gcg gtg cgc ggc ttc ggc ggc gaa gtg	384
	Thr Ala Asp Ile Lys Val Asp Ala Val Arg Gly Phe Gly Gly Glu Val	
	115 120 125	
5	ctg ctc cac ggc gcg aac ttt gat gaa gcg aaa gcc aaa gcg atc gaa	432
	Leu Leu His Gly Ala Asn Phe Asp Glu Ala Lys Ala Lys Ala Ile Glu	
	130 135 140	
10	ctg tca cag cag cag ggg ttc acc tgg gtg ccg ccg ttc gac cat ccg	480
	Leu Ser Gln Gln Gln Gly Phe Thr Trp Val Pro Pro Phe Asp His Pro	
	145 150 155 160	
15	atg gtg att gcc ggc caa ggc acg ctg gcg ctg gaa ctg ctc cag cag	528
	Met Val Ile Ala Gly Gln Gly Thr Leu Ala Leu Glu Leu Leu Gln Gln	
	165 170 175	
20	gac gcc cat ctc gac cgc gta ttt gtg cca gtc ggc ggc ggc ggt ctg	576
	Asp Ala His Leu Asp Arg Val Phe Val Pro Val Gly Gly Gly Gly Leu	
	180 185 190	
25	gct gct ggc gtg gcg gtg ctg atc aaa caa ctg atg ccg caa atc aaa	624
	Ala Ala Gly Val Ala Val Leu Ile Lys Gln Leu Met Pro Gln Ile Lys	
	195 200 205	
30	gtg atc gcc gta gaa gcg gaa gac tcc gcc tgc ctg aaa gca gcg ctg	672
	Val Ile Ala Val Glu Ala Glu Asp Ser Ala Cys Leu Lys Ala Ala Leu	
	210 215 220	
35	gat gcg ggt cat ccg gtt gat ctg ccg cgc gta ggc cta ttt gct gaa	720
	Asp Ala Gly His Pro Val Asp Leu Pro Arg Val Gly Leu Phe Ala Glu	
	225 230 235 240	
40	ggc gta gcg gta aaa cgc atc ggt gac gaa acc ttc cgt tta tgc cag	768
	Gly Val Ala Val Lys Arg Ile Gly Asp Glu Thr Phe Arg Leu Cys Gln	
	245 250 255	
45	gag tat ctc gac gac atc atc acc gtc gat agc gat gcg atc tgt gcg	816
	Glu Tyr Leu Asp Asp Ile Ile Thr Val Asp Ser Asp Ala Ile Cys Ala	
	260 265 270	
50	gcg atg aag gat tta ttc gaa gat gtg cgc gcg gtg gcg aaa ccc tct	864
	Ala Met Lys Asp Leu Phe Glu Asp Val Arg Ala Val Ala Lys Pro Ser	
	275 280 285	
55	ggc gcg ctg gcg ctg gcg gga atg aaa aaa tat atc gcc ctg cac aac	912
	Gly Ala Leu Ala Leu Ala Gly Met Lys Lys Tyr Ile Ala Leu His Asn	
	290 295 300	
60	att cgc ggc gaa cgg ctg gcg cat att ctt tcc ggt gcc aac gtg aac	960
	Ile Arg Gly Glu Arg Leu Ala His Ile Leu Ser Gly Ala Asn Val Asn	
	305 310 315 320	
65	ttc cac ggc ctg cgc tac gtc tca gaa cgc tgc gaa ctg ggc gaa cag	1008
	Phe His Gly Leu Arg Tyr Val Ser Glu Arg Cys Glu Leu Gly Glu Gln	
	325 330 335	
70	cgt gaa gcg ttg ttg gcg gtg acc att ccg gaa gaa aaa ggc agc ttc	1056
	Arg Glu Ala Leu Leu Ala Val Thr Ile Pro Glu Glu Lys Gly Ser Phe	
	340 345 350	

11

	ctc	aaa	ttc	tgc	caa	ctg	ctt	ggc	ggg	cgt	tcg	gtc	acc	gag	ttc	aac	1104
	Leu	Lys	Phe	Cys	Gln	Leu	Leu	Gly	Gly	Arg	Ser	Val	Thr	Glu	Phe	Asn	
			355					360					365				
5	tac	cgt	ttt	gcc	gat	gcc	aaa	aac	gcc	tgc	atc	ttt	gtc	ggg	gtg	cgc	1152
	Tyr	Arg	Phe	Ala	Asp	Ala	Lys	Asn	Ala	Cys	Ile	Phe	Val	Gly	Val	Arg	
		370					375					380					
10	ctg	agc	cgc	ggc	ctc	gaa	gag	cgc	aaa	gaa	att	ttg	cag	atg	ctc	aac	1200
	Leu	Ser	Arg	Gly	Leu	Glu	Glu	Arg	Lys	Glu	Ile	Leu	Gln	Met	Leu	Asn	
	385					390					395					400	
15	gac	ggc	ggc	tac	agc	gtg	gtt	gat	ctc	tcc	gac	gac	gaa	atg	gcg	aag	1248
	Asp	Gly	Gly	Tyr	Ser	Val	Val	Asp	Leu	Ser	Asp	Asp	Glu	Met	Ala	Lys	
				405						410					415		
20	cta	cac	gtg	cgc	tat	atg	gtc	ggc	gga	cgt	cca	tcg	cat	ccg	ttg	cag	1296
	Leu	His	Val	Arg	Tyr	Met	Val	Gly	Gly	Arg	Pro	Ser	His	Pro	Leu	Gln	
				420					425					430			
	gaa	cgc	ctc	tac	agc	ttc	gaa	ttc	ccg	gaa	tca	ccg	ggc	gcg	ctg	ctg	1344
	Glu	Arg	Leu	Tyr	Ser	Phe	Glu	Phe	Pro	Glu	Ser	Pro	Gly	Ala	Leu	Leu	
			435					440					445				
25	cgc	ttc	ctc	aac	acg	ctg	ggg	acg	tac	tgg	aac	att	tct	ttg	ttc	cac	1392
	Arg	Phe	Leu	Asn	Thr	Leu	Gly	Thr	Tyr	Trp	Asn	Ile	Ser	Leu	Phe	His	
		450					455					460					
30	tat	cgc	agc	cat	ggc	acc	gac	tac	ggg	cgc	gta	ctg	gcg	gcg	ttc	gaa	1440
	Tyr	Arg	Ser	His	Gly	Thr	Asp	Tyr	Gly	Arg	Val	Leu	Ala	Ala	Phe	Glu	
	465					470					475					480	
35	ctt	ggc	gac	cat	gaa	ccg	gat	ttc	gaa	acc	cgg	ctg	aat	gag	ctg	ggc	1488
	Leu	Gly	Asp	His	Glu	Pro	Asp	Phe	Glu	Thr	Arg	Leu	Asn	Glu	Leu	Gly	
				485						490					495		
40	tac	gat	tgc	cac	gac	gaa	acc	aat	aac	ccg	gcg	ttc	agg	ttc	ttt	ttg	1536
	Tyr	Asp	Cys	His	Asp	Glu	Thr	Asn	Asn	Pro	Ala	Phe	Arg	Phe	Phe	Leu	
				500					505					510			
	gcg	ggg	tag														1545
	Ala	Gly															
45	<210>	8															
	<211>	514															
	<212>	PRT															
	<213>	Escherichia coli															
50	<400>	8															
	Met	Ala	Asp	Ser	Gln	Pro	Leu	Ser	Gly	Ala	Pro	Glu	Gly	Ala	Glu	Tyr	
	1				5					10					15		
55	Leu	Arg	Ala	Val	Leu	Arg	Ala	Pro	Val	Tyr	Glu	Ala	Ala	Gln	Val	Thr	
				20					25					30			
	Pro	Leu	Gln	Lys	Met	Glu	Lys	Leu	Ser	Ser	Arg	Leu	Asp	Asn	Val	Ile	
			35					40					45				
60	Leu	Val	Lys	Arg	Glu	Asp	Arg	Gln	Pro	Val	His	Ser	Phe	Lys	Leu	Arg	
		50					55					60					

12

	Gly	Ala	Tyr	Ala	Met	Met	Ala	Gly	Leu	Thr	Glu	Glu	Gln	Lys	Ala	His	
	65					70					75					80	
5	Gly	Val	Ile	Thr	Ala	Ser	Ala	Gly	Asn	His	Ala	Gln	Gly	Val	Ala	Phe	
					85				90						95		
	Ser	Ser	Ala	Arg	Leu	Gly	Val	Lys	Ala	Leu	Ile	Val	Met	Pro	Thr	Ala	
				100				105						110			
10	Thr	Ala	Asp	Ile	Lys	Val	Asp	Ala	Val	Arg	Gly	Phe	Gly	Gly	Glu	Val	
			115					120					125				
	Leu	Leu	His	Gly	Ala	Asn	Phe	Asp	Glu	Ala	Lys	Ala	Lys	Ala	Ile	Glu	
15		130					135					140					
	Leu	Ser	Gln	Gln	Gln	Gly	Phe	Thr	Trp	Val	Pro	Pro	Phe	Asp	His	Pro	
	145					150					155					160	
20	Met	Val	Ile	Ala	Gly	Gln	Gly	Thr	Leu	Ala	Leu	Glu	Leu	Leu	Gln	Gln	
					165					170					175		
	Asp	Ala	His	Leu	Asp	Arg	Val	Phe	Val	Pro	Val	Gly	Gly	Gly	Gly	Leu	
				180					185					190			
25																	
	Ala	Ala	Gly	Val	Ala	Val	Leu	Ile	Lys	Gln	Leu	Met	Pro	Gln	Ile	Lys	
			195					200					205				
30	Val	Ile	Ala	Val	Glu	Ala	Glu	Asp	Ser	Ala	Cys	Leu	Lys	Ala	Ala	Leu	
		210					215					220					
	Asp	Ala	Gly	His	Pro	Val	Asp	Leu	Pro	Arg	Val	Gly	Leu	Phe	Ala	Glu	
	225					230					235					240	
35																	
	Gly	Val	Ala	Val	Lys	Arg	Ile	Gly	Asp	Glu	Thr	Phe	Arg	Leu	Cys	Gln	
					245					250					255		
	Glu	Tyr	Leu	Asp	Asp	Ile	Ile	Thr	Val	Asp	Ser	Asp	Ala	Ile	Cys	Ala	
40				260					265					270			
	Ala	Met	Lys	Asp	Leu	Phe	Glu	Asp	Val	Arg	Ala	Val	Ala	Lys	Pro	Ser	
			275					280					285				
45																	
	Gly	Ala	Leu	Ala	Leu	Ala	Gly	Met	Lys	Lys	Tyr	Ile	Ala	Leu	His	Asn	
		290					295					300					
	Ile	Arg	Gly	Glu	Arg	Leu	Ala	His	Ile	Leu	Ser	Gly	Ala	Asn	Val	Asn	
	305					310					315					320	
50																	
	Phe	His	Gly	Leu	Arg	Tyr	Val	Ser	Glu	Arg	Cys	Glu	Leu	Gly	Glu	Gln	
					325					330					335		
	Arg	Glu	Ala	Leu	Leu	Ala	Val	Thr	Ile	Pro	Glu	Glu	Lys	Gly	Ser	Phe	
55				340					345					350			
	Leu	Lys	Phe	Cys	Gln	Leu	Leu	Gly	Gly	Arg	Ser	Val	Thr	Glu	Phe	Asn	
			355					360					365				
60																	
	Tyr	Arg	Phe	Ala	Asp	Ala	Lys	Asn	Ala	Cys	Ile	Phe	Val	Gly	Val	Arg	
		370					375					380					

13

Leu Ser Arg Gly Leu Glu Glu Arg Lys Glu Ile Leu Gln Met Leu Asn
 385 390 395 400
 5 Asp Gly Gly Tyr Ser Val Val Asp Leu Ser Asp Asp Glu Met Ala Lys
 405 410 415
 Leu His Val Arg Tyr Met Val Gly Gly Arg Pro Ser His Pro Leu Gln
 420 425 430
 10 Glu Arg Leu Tyr Ser Phe Glu Phe Pro Glu Ser Pro Gly Ala Leu Leu
 435 440 445
 Arg Phe Leu Asn Thr Leu Gly Thr Tyr Trp Asn Ile Ser Leu Phe His
 450 455 460
 15 Tyr Arg Ser His Gly Thr Asp Tyr Gly Arg Val Leu Ala Ala Phe Glu
 465 470 475 480
 Leu Gly Asp His Glu Pro Asp Phe Glu Thr Arg Leu Asn Glu Leu Gly
 485 490 495
 Tyr Asp Cys His Asp Glu Thr Asn Asn Pro Ala Phe Arg Phe Phe Leu
 500 505 510
 25 Ala Gly
 30 <210> 9
 <211> 1548
 <212> DNA
 <213> Escherichia coli
 35 <220>
 <221> DNA
 <222> (1)..(1548)
 <223>
 40 <220>
 <221> CDS
 <222> (527)..(952)
 <223> yjgF orf
 45 <400> 9
 tcgcgatctg gtactgtaag gggaaataga gatgacacac gataataaat tgcaggttga 60
 agctattaaa cgcggcacgg taattgacca tatccccgcc cagatcgggtt ttaagctgtt 120
 50 gagtctgttc aagctgaccg aaacggatca gcgcatacacc attggtctga acctgccttc 180
 tggcgagatg ggccgcaaag atctgatcaa aatcgaaaat acctttttga gtgaagatca 240
 55 agtagatcaa ctggcattgt atgcgcgcga agccacgggtt aaccgtatcg acaactatga 300
 agtggtgggt aaatcgcgcc caagtctgcc ggagcgcata gacaatgtgc tgggtctgcc 360
 gaacagcaac tgtatcagcc atgccgaacc ggtttcatcc agctttgccg tgcgaaaacg 420
 60 cgccaatgat atcgcgctca aatgcaaata ctgtgaaaaa gagttttccc ataatgtggt 480

14

	gctggccaat taattgcggt tggtaataaa agtctggctc cctata atg agc cag	535
	Met Ser Gln	
	1	
5	act ttt tac cgc tgt aat aaa gga gaa atc atg agc aaa act atc gcg	583
	Thr Phe Tyr Arg Cys Asn Lys Gly Glu Ile Met Ser Lys Thr Ile Ala	
	5 10 15	
10	acg gaa aat gca ccg gca gct atc ggt cct tac gta cag ggc gtt gat	631
	Thr Glu Asn Ala Pro Ala Ala Ile Gly Pro Tyr Val Gln Gly Val Asp	
	20 25 30 35	
15	ctg ggc aat atg atc atc acc tcc ggt cag atc ccg gta aat ccg aaa	679
	Leu Gly Asn Met Ile Ile Thr Ser Gly Gln Ile Pro Val Asn Pro Lys	
	40 45 50	
20	acg ggc gaa gta ccg gca gac gtc gct gca cag gca cgt cag tcg ctg	727
	Thr Gly Glu Val Pro Ala Asp Val Ala Ala Gln Ala Arg Gln Ser Leu	
	55 60 65	
25	gat aac gta aaa gcg atc gtc gaa gcc gct ggc ctg aaa gtg ggc gac	775
	Asp Asn Val Lys Ala Ile Val Glu Ala Ala Gly Leu Lys Val Gly Asp	
	70 75 80	
30	atc gtt aaa act acc gtg ttt gta aaa gat ctg aac gac ttc gca acc	823
	Ile Val Lys Thr Thr Val Phe Val Lys Asp Leu Asn Asp Phe Ala Thr	
	85 90 95	
35	gta aac gcc act tac gaa gcc ttc ttc acc gaa cac aac gcc acc ttc	871
	Val Asn Ala Thr Tyr Glu Ala Phe Phe Thr Glu His Asn Ala Thr Phe	
	100 105 110 115	
40	ccg gca cgt tct tgc gtt gaa gtt gcc cgt ctg ccg aaa gac gtg aag	919
	Pro Ala Arg Ser Cys Val Glu Val Ala Arg Leu Pro Lys Asp Val Lys	
	120 125 130	
45	att gag atc gaa gcg atc gct gtt cgt cgc taa tcttgatgga aatccgggct	972
	Ile Glu Ile Glu Ala Ile Ala Val Arg Arg	
	135 140	
50	atcatgcccg gattaagtct gatgacaaac gcaaaatcgc ctgatgcgct acgcttatca	1032
	ggcctacgtg attcctgcaa tttattgaat ttgttggccg gataaggcat ttacgccgca	1092
55	tccggcatga acaaaaactca ctttgtctac aatctgaatc ggggctatcg tgcccagttt	1152
	attctttatt gccagccgta acgacggcta tagaaccctt tcaccaactg ggttaatgtc	1212
60	atataccctg ccagaatcgc aaccagccac gggaaatagc ttaacggcag cgcctgtaat	1272
	tgcagataac tggccagcgg tga aaacggc aatgcgatcc cgacaatcat cagatcacg	1332
	gtcatgatca ttaacggcca cgatgcacag ctctgaataa acggcacacg gcgggtgcgg	1392
	atcatatgca caatcagcgt ttgcgacagt aagcccacca caaaccatcc cgactggaac	1452
	agcgttttgcg tttccggcgt gttggcatgg aatacccacc acatcaggca aaacgtcaaa	1512
	atatcgaaga tcgagctgat cggtccgaag aagatc	1548

15

<210> 10
 <211> 141
 <212> PRT
 <213> Escherichia coli
 5

<400> 10
 Met Ser Gln Thr Phe Tyr Arg Cys Asn Lys Gly Glu Ile Met Ser Lys
 1 5 10 15
 10 Thr Ile Ala Thr Glu Asn Ala Pro Ala Ala Ile Gly Pro Tyr Val Gln
 20 25 30
 15 Gly Val Asp Leu Gly Asn Met Ile Ile Thr Ser Gly Gln Ile Pro Val
 35 40 45
 Asn Pro Lys Thr Gly Glu Val Pro Ala Asp Val Ala Ala Gln Ala Arg
 50 55 60
 20 Gln Ser Leu Asp Asn Val Lys Ala Ile Val Glu Ala Ala Gly Leu Lys
 65 70 75 80
 Val Gly Asp Ile Val Lys Thr Thr Val Phe Val Lys Asp Leu Asn Asp
 85 90 95
 25 Phe Ala Thr Val Asn Ala Thr Tyr Glu Ala Phe Phe Thr Glu His Asn
 100 105 110
 Ala Thr Phe Pro Ala Arg Ser Cys Val Glu Val Ala Arg Leu Pro Lys
 115 120 125
 30 Asp Val Lys Ile Glu Ile Glu Ala Ile Ala Val Arg Arg
 130 135 140

INTERNATIONAL SEARCH REPORT

 International Application No
 PCT/EP2004/008389

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12P13/08 C12R1/19 //(C12P13/08,C12R1:19)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12P C12R

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, BIOSIS, FSTA, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DEBABOV V G: "The threonine story" ADVANCES IN BIOCHEMICAL ENGINEERING, BIOTECHNOLOGY, SPRINGER, BERLIN, DE, vol. 79, 2003, pages 113-136, XP008014933 ISSN: 0724-6145	35
Y	the whole document figure 2 page 124, line 12-27 ---	1-39
Y	DEBABOV V: "Construction of strains producing L-threonine" PROCEEDINGS OF THE INTERNATIONAL SYMPOSIUM ON THE GENETICS OF INDUSTRIAL MICROORGANISMS, TOKYO, JAPAN, 1982, pages 254-258, XP008031179 cited in the application the whole document --- -/--	1-39

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *Z* document member of the same patent family

Date of the actual completion of the international search

8 December 2004

Date of mailing of the international search report

28/12/2004

Name and mailing address of the ISA

 European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

van de Kamp, M

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP2004/008389

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	<p>SHIU-LAN L ET AL: "Study on the microfiltration of Escherichia coli-containing fermentation broth by a ceramic membrane filter"</p> <p>JOURNAL OF MEMBRANE SCIENCE, vol. 110, no. 2, 21 February 1996 (1996-02-21), pages 203-210, XP004041728</p> <p>ISSN: 0376-7388</p> <p>abstract</p> <p>page 203, left-hand column, line 1</p> <p>-right-hand column, line 16</p> <p>figure 2</p> <p>---</p>	1-34, 36-39
Y	<p>TODA KIYOSHI: "Theoretical and methodological studies of continuous microbial bioreactors."</p> <p>JOURNAL OF GENERAL AND APPLIED MICROBIOLOGY, vol. 49, no. 4, August 2003 (2003-08), pages 219-233, XP008038115</p> <p>ISSN: 0022-1260</p> <p>cited in the application</p> <p>abstract</p> <p>page 228, right-hand column, line 38 -page 230, left-hand column, line 11</p> <p>table 2</p> <p>page 230, left-hand column, line 12-41</p> <p>---</p>	1-34, 36-39
Y	<p>ZENG A-P: "Continuous culture" 1999, IN: "MANUAL OF INDUSTRIAL MICROBIOLOGY AND BIOTECHNOLOGY" (ED. DEMAÏN A L ET AL), 2ND ED., ASM PRESS, WASHINGTON XP002305534</p> <p>ISBN: 1-55581-128-0</p> <p>page 151-164</p> <p>figures 1C,8</p> <p>paragraphs 12.1.4, 12.2.1 and 12.2.2.5</p> <p>---</p>	1-34, 36-39
A	<p>US 6 025 169 A (KOYAMA YOSUKE ET AL) 15 February 2000 (2000-02-15)</p> <p>cited in the application</p> <p>figures 1,2</p> <p>examples 1,4,6-9</p> <p>claims 1-3</p> <p>---</p>	1
A	<p>EP 0 139 592 A (SANTERRE ORSAN) 2 May 1985 (1985-05-02)</p> <p>cited in the application</p> <p>examples 4,5,7-12,15</p> <p>claims 1,2,6,7,9,10</p> <p>figures 2,5</p> <p>---</p> <p style="text-align: center;">-/--</p>	1

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP2004/008389

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 5 260 216 A (HIROSE TOSHIKI ET AL) 9 November 1993 (1993-11-09) cited in the application figure 1 column 2, line 1-34 example 2 claim 1 ---	1
A	OKAMOTO K ET AL: "Development of an industrially stable process for L-threonine fermentation by an L-methionine-auxotrophic mutant of Escherichia coli" JOURNAL OF BIOSCIENCE AND BIOENGINEERING, vol. 89, no. 1, 6 January 2000 (2000-01-06), pages 87-89, XP002305525 ISSN: 1389-1723 abstract ---	1
A	OKAMOTO K ET AL: "Hyperproduction of L-threonine by an Escherichia coli mutant with impaired L-threonine uptake" BIOSCIENCE BIOTECHNOLOGY BIOCHEMISTRY, vol. 61, no. 11, November 1997 (1997-11), pages 1877-1882, XP001018682 ISSN: 0916-8451 cited in the application abstract ---	1
A	OBON J M ET AL: "High-density Escherichia coli cultures for continuous L(-)-carnitine production" APPLIED MICROBIOLOGY AND BIOTECHNOLOGY, vol. 51, no. 6, June 1999 (1999-06), pages 760-764, XP002305526 ISSN: 0175-7598 abstract figure 1 ---	1
A	CANOVAS M ET AL: "Modeling of the biotransformation of crotonobetaine into L-(-)-carnitine by Escherichia coli strains" BIOTECHNOLOGY AND BIOENGINEERING, vol. 77, no. 7, 30 March 2002 (2002-03-30), pages 764-775, XP002305527 ISSN: 0006-3592 abstract --- -/--	1

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP2004/008389

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>ALVAREZ-VASQUEZ F ET AL: "Modeling, optimization and experimental assessment of continuous L-(-)-carnitine production by Escherichia coli cultures." BIOTECHNOLOGY AND BIOENGINEERING. 30 DEC 2002, vol. 80, no. 7, 30 December 2002 (2002-12-30), pages 794-805, XP002305528 ISSN: 0006-3592 abstract</p> <p>---</p>	1
A	<p>YU PENG ET AL: "Continuous production of cell-free recombinant proteins using Escherichia coli" BIOTECHNOLOGY PROGRESS, vol. 9, no. 6, 1993, pages 587-593, XP002305529 ISSN: 8756-7938 abstract figure 1</p> <p>---</p>	1
A	<p>HENRY K L ET AL: "Continuous recombinant bacterial fermentations utilizing selective flocculation and recycle." BIOTECHNOLOGY PROGRESS. 1990 JAN-FEB, vol. 6, no. 1, January 1990 (1990-01), pages 7-12, XP002305530 ISSN: 8756-7938 abstract figure 1</p> <p>---</p>	1
A	<p>ZENG A-P ET AL: "Production of 2,3-butanediol in a membrane bioreactor with cell recycle" APPLIED MICROBIOLOGY AND BIOTECHNOLOGY, vol. 34, no. 4, 1991, pages 463-468, XP008038261 ISSN: 0175-7598 abstract figure 1</p> <p>---</p>	1
A	<p>BULL D N ET AL: "Enhanced product formation in continuous fermentations with microbial cell recycle" BIOTECHNOLOGY AND BIOENGINEERING, vol. 23, no. 2, 1981, pages 373-389, XP002305532 ISSN: 0006-3592 abstract figures 1,7</p> <p>---</p> <p style="text-align: center;">-/--</p>	1

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP2004/008389

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	BRYERS J D: "Effects of cell recycle on microbial cell viability and metabolism" JOURNAL OF PARENTERAL SCIENCE AND TECHNOLOGY, vol. 42, no. 2, 1988, pages 40-46, XP008038268 ISSN: 0279-7976 abstract figures 2-4 ---	1
A	SCHUGERL K: "Integrated processing of biotechnology products" BIOTECHNOLOGY ADVANCES, ELSEVIER PUBLISHING, BARKING, GB, vol. 18, no. 7, 1 November 2000 (2000-11-01), pages 581-599, XP004317587 ISSN: 0734-9750 abstract ---	1
A	VAN DER WIELEN L A M ET AL: "Integrated product formation and recovery in fermentation" CURRENT OPINION IN BIOTECHNOLOGY, vol. 3, no. 2, 1992, pages 130-138, XP002304966 ISSN: 0958-1669 abstract table 2 figure 2 -----	1

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP2004/008389

Patent document cited in search report		Publication date		Patent family member(s)	Publication date
US 6025169	A	15-02-2000	JP	3074781 B2	07-08-2000
			JP	5030985 A	09-02-1993
			JP	2932791 B2	09-08-1999
			JP	5076346 A	30-03-1993
			US	5912113 A	15-06-1999
			BE	1008008 A3	12-12-1995
			BR	9105208 A	21-07-1992
			FR	2669935 A1	05-06-1992
			FR	2676234 A1	13-11-1992
			IT	1256566 B	11-12-1995
			CN	1064890 A , B	30-09-1992
			CN	1183474 A , B	03-06-1998
EP 0139592	A	02-05-1985	FR	2554126 A1	03-05-1985
			AT	74157 T	15-04-1992
			DE	3485613 D1	30-04-1992
			DE	139592 T1	22-05-1986
			EP	0139592 A2	02-05-1985
			ES	8506805 A1	16-11-1985
			JP	60133891 A	17-07-1985
			KR	9301263 B1	22-02-1993
US 5260216	A	09-11-1993	JP	3018449 B2	13-03-2000
			JP	4131087 A	01-05-1992
			BE	1006589 A3	25-10-1994
			FR	2667076 A1	27-03-1992
			ID	1001 B	10-10-1996
			IT	1251203 B	04-05-1995
			US	5362635 A	08-11-1994